No.



200300170

HEIR UNIVERD SYNAVIES OF ANTERIOA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

Jexus Agriculturul Experiment Stution

MIRRORS, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR IMPORTING IT, OR EXPORTING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE ABOVE PURPOSE, OR CONDITIONING IT REPOPAGATION, OR STOCKING IT FOR ANY OF THE ABOVE PURPOSE, OR USING IT IN PRODUCING A HYBRID OR TRENT VARIETY THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY PROTECTION ACT. IN THE STATES SEED OF THIS VARIETY (1) SHALL BE SOLD BY VARIETY NAME ONLY AS A CLASS OF CERTIFIED SOLD SY SHALL CONFORM TO THE NUMBER OF GENERATIONS SPECIFIED BY THE OWNER OF THE RIGHTS. (84 STAT. 1542)

PEANUT

'Tamrun OL 02'

In Testimonn Moscost, I have hereunto set my hand and caused the seal of the Minni Pariety Protection Office to be affixed at the City of Washington, D.C. this tenth day of November, in the year two thousand and four.

Commissioner
Plant Variety Protection Office

of Agriculture

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY - PLANT VARIETY PROTECTION OFFICE

APPLICATION FOR PLANT VARIETY PROTECTION CERTIFICATE

(Instructions and information	collection burden statement	on reverse	e)	,				
1. NAME OF OWNER	<u> </u>				2. TEMPORARY DESIGNAT EXPERIMENTAL NAME	ION OR	3. VARIETY NAME	
Texas Agricultural	Experiment Sta	tion			Tx977053		Tamrun OL O	2
4. ADDRESS (Street and No., or R.F.D. No.	, City, State, and ZIP Code, and Cour	ntry)			5. TELEPHONE (include area	a code)	FOR OFFICIAL USE ONL	Υ
Dr. Frank E. Gilst	rap				979-845-4747		PVPO NUMBER	
Associate Agency I 2147 TAMU	Director				6. FAX (include area code)	2	003001	70
College Station, T	rx 77843-2147			 	979-458-4765		FILING DATE	
7. IF THE OWNER NAMED IS NOT A "PERSORGANIZATION (corporation, partnership State of Texas Rese	SON", GIVE FORM OF b, association, etc.) Barch Agency	8. IF INCO	ORPORATED, GIVE OF INCORPORA	VE TION	9. DATE OF INCORPORATION	ON	February	25, 7
Janie Hurley Technology Licensi Technology Licensi The Texas A&M Univ 3369 TAMU College Station, T	eresentative(s) to serve in the ing Manager, Agri ing Office versity System						FILING AND EXAMINATION FEES: S 3,652.00 DATE 2 25 03 CERTIFICATION FEE: Y 432.00 DATE 10 27	9
11. TELEPHONE (Include area code) 979-847-8682	12. FAX (Include area code) 979-845-1402	-	13. E-MAIL jhurle	y@tam	u.edu	! _	OP KIND (Common Name) eanut	
15. GENUS AND SPECIES NAME OF CROP L. subsp. <u>hypogaea</u>		aea	16. FAMILYNA Legumi	•	0	17. IS TH HYBR	HE VARIETY A FIRST GENERAT	ION
repository)	History of the Variety tness n of Variety n of the Variety (<i>Optional</i>)	ted varieties, approved pub	20. CV	OOES THE OF ARIETY BE IT OOES THE OF ARIETY BE IT OOES THE OF ARIETY BE IT OOES THE O	ES (if "yes", answer items 20 and 21 below) WNER SPECIFY THAT SEED (.IMITED AS TO NUMBER OF (H CLASSES? WNER SPECIFY THAT SEED (.IMITED AS TO NUMBER OF (.IMITED AS TO NUMBER OF (IFY THE T FOUNDATION	OF THIS CLASSES? ATION [OF THIS SENERATION]	TREGISTERED TO CERT	 NO
22. HAS THE VARIETY (INCLUDING ANY HAF FROM THIS VARIETY BEEN SOLD, DISF OTHER COUNTRIES? YES IF YES, YOU MUST PROVIDE THE DATI FOR EACH COUNTRY AND THE CIRCU	⊠ NO		ED 23, 15 : S. OR P	THE VARIE ROPERTY R		THE VARIE GHT OR PA	ETY PROTECTED BY INTELLEC ITENT)? NO I ISSUANCE AND ASSIGNED	TUAL
24. The owners declare that a viable sample of for a tuber propagated variety a tissue cult. The undersigned owner(s) is(are) the owner and is entitled to protection under the providence; is (are) informed that false represent the providence.	er of this sexually reproduced or tuber isions of Section 42 of the Plant Varie	r propagated party Protection	olant variety, and b Act.					
SIGNATURE OF OWNER	Gisto		SIGN	ATURE OF O	WNER			
NAME (Please print or type) Frank E. Gilstrap			NAME	: (Please prin	t or type)			
capacity or title Associate Agency Dir	cector, TAES 2	1410	3 CAPA	CITY OR TIT	LE		DATE	
T 470 (07 01) deciseed by the Bleet Verich, Br	rotaction Office with MovelDe deat C.O.	D1- 07	ED 470 (04 04)	*. 1. 1 1 1	· Con minare for leafer		Information antiqueling burden seek	

INSTRUCTIONS

GENERAL: To be effectively filed with the Plant Variety Protection Office (PVPO), ALL of the following items must be received in the PVPO: (1) Completed application form signed by the owner; (2) completed exhibits A, B, C, E; (3) for a seed reproduced variety at least 2,500 viable untreated seeds, for a hybrid variety at least 2,500 untreated seeds of each line necessary to reproduce the variety, or for tuber reproduced varieties verification that a viable (in the sense that it will reproduce an entire plant) tissue culture will be deposited and maintained in an approved public repository; (4) check drawn on a U.S. bank for \$2,705 (\$320 filing fee and \$2,385 examination fee), payable to "Treasurer of the United States" (See Section 97.6 of the Regulations and Rules of Practice.) Partial applications will be held in the PVPO for not more than 90 days, then returned to the applicant as unfilled. Mail application and other requirements to Plant Variety Protection Office, AMS, USDA, Room 401, NAL Building, 10301 Baltimore Avenue, Beltsville, MD 20705-2351. Retain one copy for your files. All items on the face of the application are self explanatory unless noted below. Corrections on the application form and exhibits must be initialed and dated. DO NOT use masking materials to make corrections. If a certificate is allowed, you will be requested to send a check payable to "Treasurer of the United States" in the amount of \$320 for issuance of the certificate. Certificates will be issued to owner, not licensee or agent.

Plant Variety Protection Office Telephone: (301) 504-5518 FAX: (301) 504-5291

Homepage: http://www.ams.usda.gov/science/pvpo/pvp.htm

ITEM

18a, Give:

- (1) the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method;
- (2) the details of subsequent stages of selection and multiplication;
- (3) evidence of uniformity and stability; and
- (4) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified
- 18b. Give a summary of the variety's distinctness. Clearly state how this application variety may be distinguished from all other varieties in the same crop. If the new variety is most similar to one variety or a group of related varieties:
 - (1) identify these varieties and state all differences objectively;
 - (2) attach statistical data for characters expressed numerically and demonstrate that these are clear differences; and
 - (3) submit, if helpful, seed and plant specimens or photographs (prints) of seed and plant comparisons which clearly indicate distinctness.
- 18c. Exhibit C forms are available from the PVPO Office for most crops; specify crop kind. Fill in Exhibit C (Objective Description of Variety) form as completely as possible to describe your variety.
- 18d. Optional additional characteristics and/or photographs. Describe any additional characteristics that cannot be accurately conveyed in Exhibit C. Use comparative varieties as is necessary to reveal more accurately the characteristics that are difficult to describe, such as plant habit, plant color, disease resistance, etc.
- 18e. Section 52(5) of the Act requires applicants to furnish a statement of the basis of the applicant's ownership. An Exhibit E form is available from the PVPO.
- 19. If "Yes" is specified (seed of this variety be sold by variety name only, as a class of certified seed), the applicant MAY NOT reverse this affirmative decision after the variety has been sold and so labeled, the decision published, or the certificate issued. However, if "No" has been specified, the applicant may change the choice. (See Regulations and Rules of Practice, Section 97.103).
- 22. See Sections 41, 42, and 43 of the Act and Section 97.5 of the regulations for eligibility requirements.
- 23. See Section 55 of the Act for instructions on claiming the benefit of an earlier filing date.
- 21. CONTINUED FROM FRONT (Please provide a statement as to the limitation and sequence of generations that may be certified.)

22. CONTINUED FROM FRONT— (Please provide the date of first sale, disposition, transfer, or use for each country and the circumstances, if the variety (including any harvested material) or a hybrid produced from this variety has been sold, disposed of, transferred, or used in the U.S. or other countries.)

- 23. CONTINUED FROM FRONT (Please give the country, date of filing or issuance, and assigned reference number, if the variety or any component of the variety is protected by intellectual property right (Plant Breeder's Right or Patent).)
 - A component of this variety is protected, per a license agreement with the University of Florida Research Foundation; U.S. Patent Nos. 5,922,390; 6,063,984; 6,121,472.

NOTES: It is the responsibility of the applicant/owner to keep the PVPO informed of any changes of address or change of ownership or assignment or owner's representative during the life of the application/certificate. There is no charge for filing a change of address. The fee for filing a change of ownership or assignment or any modification of owner's name is specified in Section 97.175 of the regulations. (See Section 101 of the Act, and Sections 97.130, 97.131, 97.175(h) of the Regulations and Rules of Practice.)

To avoid conflict with other variety names in use, the applicant must check the appropriate recognized authority. For example, for agricultural and vegetable crops, contact: Seed Branch, AMS, USDA, Room 213, Building 306, Beltsville Agricultural Research Center--East, Beltsville, MD 20705. Telephone: (301) 504-8089. http://www.ams.usda.gov/lsg/seed.htm

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 3.0 hours per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, sexual orientation, or marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audictape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

S&T-470 (07-01) designed by the Plant Variety Protection Office with WordPerfect 9.0. Replaces STD-470 (04-01) which is obsolete.

Exhibit A.

ORIGIN AND BREEDING HISTORY OF 'TAMRUN OL 02'

'Tamrun OL 02,' tested under experimental number Tx977053, was derived from a backcross between Tx896100 and 'Sun Oleic 95R' that was made at College Station, Texas in 1996. Tx896100 was later released as 'Tamrun 96' because it had excellent yield potential under heavy Tomato Spotted Wilt Virus (TSWV) pressure and had reduced incidence of pod rot and southern blight. Since the release of Tamrun 96 it has become widely grown in areas of Sclerotinia blight infection because of its ability to yield under moderate sclerotinia disease pressure. The disease resistance and yield potential of Tamrun 96 were the traits that attracted the TAMU Breeding Program to initially choose this line as the recurrent parent in introducing the much sought after high $O/L^{1/2}$ trait.

The BC_1F_1 seed were grown in the summer of 1996 in a field near Bryan, Texas. The BC_1F_2 generation was space planted in a winter nursery during 1996-97 in Puerto Rico in an attempt to accelerate the program and meet the industry's demand for quick high O/L variety development. Individual BC_1F_2 progeny were harvested and grown as $BC_1F_{2:3}$ plant rows in a TSWV screening nursery near Pearsall, Texas in 1997. Plant rows were selected based on disease ratings and agronomic traits. The selected lines were grown in 1998 and 1999 in two preliminary $F_{2:4}$ and $F_{2:5}$ yield trials at Pearsall. Selections were made based on disease ratings, yield, and grade characteristics. Individual seeds from the 2000 $BC_1F_{2:6}$ yield tests were tested for O/L value and then bulked and planted as $BC_1F_{2:7}$ breeder seed increase. From this increase, $BC_1F_{2:8}$ individual seeds were again tested for high O/L and any seeds testing less than 18:1 O/L were discarded. The remaining seed were increased and are being grown for a second breeder seed increase. The material being released is $BC_1F_{2:8}$ generation.

The variety 'Tamrun OL02' has been observed for 8 generations since selection, and is stable and uniform, except for variants in pod type; up to 0.05 % in a sample may be variant, with three seeded pods, not two seeded. This variant is commercially acceptable and predictable.

^{1/} It has been determined that a high oleic acid to linoleic acid ratio (O/L) in peanut (and other crops) imparts desirable traits into the peanut (and other crops) after it is roasted. Desirable attributes include longer shelf life of products, including peanut butter, peanut candies, and roasted peanuts in and out of the shell. Also, there are definite health benefits from eating such products, i.e., more mono- and poly-unsaturated fat in the diet. Two dominant genes reportedly control the trait but our research has shown there may be some modifiers involved, at least in the Spanish types (Lopez et al., 2000).

Exhibit B.

STATEMENT OF DISTINCTIVENESS OF 'TAMRUN OL 02'

'Tamrun OL 02' is a runner market-type peanut (Arachis hypogaea L. subsp. hypogaea var. hypogaea) cultivar with a high O/L ratio and good yield potential. The new variety was tested as Tx977053. Tx977053 is a sister line to Tamrun OL01 (Tx977006) but it has a smaller seed that is more typical of currently grown runner cultivars such as Tamrun 96. Tx977053 also has lower sugar content than Tamrun OL01. The objective in releasing Tx9770053 is to provide the peanut growers of Texas, and especially West Texas, a runner variety which has high O/L and some ability to withstand effects of soil borne diseases and tomato spotted wilt virus, but with a smaller seed size and lower sugar content than Tamrun OL01.

'Tamrun OL 02' is most similar to Florunner and Tamrun 96; however, Florunner and Tamrun 96 have smaller size seed weight per 100 sd. (Florunner = 59.8 g/100, Tamrun 96 = 60.5, vs. Tamrun OL 02 = 64.7 g/100), lower oleic acid: linoleic acid ratios (Florunner = 1.4:1, Tamrun 96 = 1.5:1, vs. Tamrun OL 02 = 19.1:1), and higher Iodine values (Florunner = 96.6, Tamrun 96 = 97.2, vs. Tamrun OL 02 = 76.8), than 'Tamrun OL 02.' More information is provided below.

Plant description: The plants of Tx977053 are profusely branched at or near the cotyledonary nodes at the base of the plant, like Florunner and Tamrun 96. Tx977053 is a larger growing plant than Florunner and Tamrun 96. Main stem height averaged 43.6 cm at three locations for Tx977053 (Table A), significantly taller than Florunner at 37.6 cm, Tamrun 96 at 31.5 cm and 29.4 cm for Tamrun OL 01. This difference is obvious in field plantings when the four lines are growing adjacent to one another, but less noticeable in a field of Tx977053. Lateral branch length for Tx977053 averaged longer than for Florunner (61.6 cm vs. 55.4 cm), but was not statistically significant (DMRT, 0.05). See Table A and Appendix I for plant measurement analyses. The plant color appears to be a darker green than Florunner (Tx977053 color is 146A, according to the Royal Horticultural Society color chart).

Leaf measurements show that the leaflets of Tx977053 are significantly longer and wider than those of Florunner, Tamrun 96, and Tamrun OL 01. (Table A and Appendix I).

Pod and seed description: The pods and seed of Tx977053 are significantly smaller than Tamrun OL 01 (Table B and Appendix II). Where the pods of Tamrun OL 01 will almost always grade 40% or more Fancy (Grade as a Virginia), Tx977053 rarely will grade as a Virginia type. Seed size for Tx977053 averaged 64.7 g/100 sd., whereas seed size for Florunner averaged 59.8 g/100 sd. Although Tx977053 is larger than Florunner and Tamrun 96 (Table B and Appendix II) shelling data show that the line partitions about equal in jumbo and medium sizes (Appendix III). Pod length for TX977053 was 31.4 mm and width = 11.8 mm. This is smaller than Tamrun OL 01 at 33.6 mm long and 13.2 mm wide. The Tx977053 pods are longer than Florunner (31.4 mm vs. 28.6 mm) but are equal in other measurements. The same is true for Tx977053 vs. Tamrun 96. The pods are moderately reticulated, like Florunner, Tamrun 96, and Tamrun OL01, but have a slightly more prominent beak than Florunner, like Tamrun 96 and Tamrun OL 01. Pods are mostly two seeded, with variant three seeded pods occurring at a rate of up to 0.05 % in a sample. The constriction between the two pod segments is moderate, but more pronounced than Florunner and Tamrun 96, although not significant statistically (data in Table B).

Seed of Tx977053 are smaller than Tamrun OL 01 in most respects, but measure to be wider than Tamrun OL 01 in the apical position, giving the apical seed a flat appearance, like Tamrun 96. The 100 seed weights (Table 1) reflect the smaller size (than Tamrun OL 01) but also emphasize that Tx977053 pods and seed are larger size than Florunner and Tamrun 96.

Chemical and Organoleptic evaluation: Quality analyses indicated significant differences between Tx977053 and Florunner in traits such as O/L ratio = 19.1:1, and Iodine number = 76.8, with oil content = 49.55 %, and protein content = 25.99 %, about equal to Florunner. The chemical analyses of this line are of primary importance in our decision to submit the line for release. The oleic acid: linoleic acid ratio of this line averaged 19.1:1 in the 2001 tests from three locations: Frio County, Erath County, and Gaines County, Texas (see **Table D-2**). The Leek company analyses gave us an average O/L for all samples run of 18.55. If the line is 15:1 O/L range, or above, in West Texas it will be acceptable because industry personnel (Hershey, M&M Mars, and others) assure us that if the O/L for a peanut as high as 15:1, other parts of the product (i.e., candy bar) will break down before the peanut oxidizes and takes on a bad flavor.

Note: The oleic acid is the acid designated as 18.1 in the table in Appendix III. Linoleic acid is the 18.2.

The 19.1:1 ratio of Tx977053 compares to a 1.4:1 for Florunner at the same locations, and 1.5:1 for Tamrun 96 (Table D-2). Actual percentages of oleic and linoleic acids are shown in Appendix III. These chemical and organoleptic evaluations were conducted by either the Leek Company, or the USDA lab in Raleigh, NC, or in our labs in College Station, Texas and Lubbock. The chemical tests done by the Leek Company were run in the Brownfield, TX lab and/or in the Pert Labs Division of Leek in Edenton, NC.

In other chemical analyses (**Appendix III**) Tx977053 was higher than Tamrun 96 and Tamrun OL 01 in oil percent (Tx977053 = 49.55; Tamrun 96 = 47.04; Tamrun OL 01 = 47.38) but equal to Florunner (Flo. = 50.5). In protein percent Tx977053 was essentially equal to Florunner, Tamrun 96, and Tamrun OL 01; (Tx977053 = 25.99; Florunner = 25.85; Tamrun 96 = 25.75; and Tamrun OL 01 = 25.16). The Iodine value (a measure of the oxidative properties of the oil which pertains to the shelf life of roasted product) of the Tx977053 is much better than Tamrun 96 and Florunner (Tx977053 = 76.8; Tamrun 96 = 97.2; Flo. = 96.6). Readings below 92 are desirable, and below 85 are considered exceptional in oxidative properties. We expected this in Tx977053 because Iodine number is directly related to the O/L ratio.

The sugar content of Tx977053 is another very important aspect of the decision to apply for release of this line. Tamrun OL01 has a slightly higher sugar content than Tx977053 and, thus, causes manufacturers difficulty in roasting the peanut. Tx977053 is lower in sugar content than Tamrun OL01, as well as Flavor Runner 458 (Table E). This character will make the line even more desirable for growers and users alike. Additional sugar content data are shown in **Appendix III**, verifying our findings.

Taste panel evaluations were very positive for the Tx977053 line. The characters evaluated included Roasted Peanutty, Sweet Aromatic, Sweet Taste, Raw Beany Green, Dark Roast, and Woody Hull Skins. Also measured was Bitterness and Astringency. The industry definitions of these characters are shown in Appendix III. Averages for these characters were all within the acceptable range for Tx977053. Analyses and exact figures for these taste tests are shown in **Appendix III**.

Shelling properties: Figures 1 and 2 (in Appendix III) show the relative shelling properties of Tx977053, Florunner, Tamrun 96, and Tamrun OL 01, indicating that the Tx977053 line is very similar to the Florunner and Tamrun 96 lines, and that these three have a very different turnout of Jumbo, Medium, No. 1's, -16 (less than 16, called other kernels), and splits than does Tamrun OL 01. From Figures 1, Figure 2, and Table 1 we see that all the characters measured in shelling were different from Tamrun OL 01, including the percent of Jumbo kernels which is significantly higher (see Appendix III) for Tamrun OL 01 than for Tx977053, Florunner, and Tamrun 96. Supportive shelling properties data are shown in Appendix III.

In measurements of bulk density of the pods of our samples, Tx977053 was in the same range as Tamrun 96 and Florunner. These measurements relate to storage properties of the line. Supportive data are shown in **Appendix III.**

Table D-2: Oleic/linoleic acid ratios and percentages

Analyses of samples from the 2001 test year

Three locations:

Frio County in South Texas Erath County in Central Texas Gaines County in West Texas

	Tamrun OL02	Tamrun 96	Florunner
Frio County	19.8* (79.3:4.0)**	1.7 (50.3:29.4)	1.6 (48.8:31.1)
Erath County	18.0 (79.7:4.0)	1.4 (46.6:33.3)	1.3 (44.7:35.4)
Gaines County	19.5 (80.1:4.1)	1.4 (45.9:33.3)	1.4 (47.5:32.9)
Variety mean for O/L	19.1	1.5	1.4

^{*}O/L ratio shown first (i.e., 19.8 to 1, oleic to linoleic acid ratio)

^{**} Percent of oleic acid and percent linoleic acid shown in parenthesis

OBJECTIVE DESCRIPTION OF VARIETY

PEANUT (Arachis hypogaea) subsp. hypogaea var. hypogaea

NAME OF APPLICANTIS	Typogaea var. nypogaea
Texas Agricultural Experiment Station	DESIGNATION
ADDRESS (Street and No. or R.F.D. No., City, State, and ZIP Code)	Tx 977053
1229 North US Hwy 281 Williams Administration Bldg Stephenville, TX 76401 Texas A&N University	FOR OFFICIAL USE ONLY
College Station, TX 77843	200300170
Place the appropriate number that describes the varietal character of this variety in th	a bassa balasi
Place a zero in first box (e-s- 0 8 9 or 0 9) when number is either 99 or less o	r 9 or less.
1. BOTANICAL TYPE:	
1 - ABSENT 2 - PRESENT	
1 = ALTERNATE — Pairs of vegetative & reproductive branches (Virgin Branching Pattern: 2 = SEQUENTIAL — Continuous reproductive branches (Valencia—Span	sia) 3- OTHER (Specify) Alternate but nish) not perfectly so.
2. PLANT:	
1 1 Di Branching:	= SPARSE (Valencia) 2 = MODERATE (Starr) 3 = PROFUSE (Florunner)
3. MATURITY:	
3 Region: 1=VIRGINIA, NORTH CAROLINA 2 = S.E. UNITED STATES 3=S.V	Y.UNITED STATES 4-OTHER
1 4 5 NUMBER OF DAYS TO MATURITY	3
0 0 NO. OF DAYS EARLIER THAN	2 = FLORUNNER 3 = FLORIGIANT 51R 5 = NC - 2
0 5 NO. OF DAYS LATER THAN	7 = SOUTHEASTERN RUNNER 56-15 ecify) Tamrun 96
LEAVES:	
COLOR AT 60 DAYS: (Nickerson Color Designation): 1 = LIGHT GREEN (10Gy 6A) 3 = DARK GREEN (5G 4/7)	2 = MEDIUM GREEN (2.5G 5/9) 4 = OTHER (Specify) 146A RHS Chart
6 0 MM. LEAFLET LENGTH (Basal leaflet of the youngest fully opened leaf)	
2.7 2 LEAFLET LENGTH/WIDTH RATIO	
POD: (Average for 20 pods at maturity)	
3 1 MM. LENGTH	METER
5 2 2 4 KG./HA. POD YIELD	
0 2 1 % LESS THAN	2 = FLORUNNER 3 = FLORIGIANT
0 0 6 * MORE THAN	18 5 = NC · 2 7 = SOUTHEASTERN RUNNER 56-15 cify) Tampun OF OT
NA % FANCY SIZE: (% riding 13.46 mm., 34/64 inch, spacing set on presizer roller)	

,,	for 20 pods at matur ER OF SEEDS PER I		2 3-3	4 - 3 -4	5 * 2 -3-4) 			
1	RICTION: 1 - SHA	LLOW OR NONE(l'irgi	niu S6R, Arge	ntine)	2 - MEDIŲI	M (Virginia 61K)	3 • DE	EP (Starr)	
1 SURFA	CE: 1 = GL/	ABROUS (Florunner)	2 - PU8	ESCENT	(Florispan)				
BEAK:	1 - A85	SENT 2 = 1	NCONSPICU	ous	3 = PRONO	UNCED			
6. SEED (Mature	, cured but not aged):							
0 3 004	T COLOR: 6 - 1	WHITE (<i>Pearl</i>) 2 = 0 RED 7 = P OTHER (<i>Specify</i>)	PURPLE 8	= TAN (S	terr) 4 = BRO PURPLE		((Florigian))	
2 COATS	SURFACE: 1 = S	SMOOTH -2 = IND	ENTED	2	1 = UNIFORM	COLOR 2	- BLEMISH	ED	
3 SHAPE		. (Starr) 2 = SHO							
1 6 MM.	LENGTH	0 9 MM, WIDTI	6	4	GRAMS PER 1	00 SEED (8% Md	oisture)		
7. DISEASE RES	ISTANCE: (O = No	t Tested, 1 = Susceptible	, 2 = Resistan	t)					
1 ѕоитн	ERN STEM ROT			1 4	UST				
1 EARLY	LEAF SPOT			0 4	IRUS X				
] SOUTH	ERN LEAF SPOT			0 "	IOSAIC				
	T COMPLEX			0 0	THER (Specify)				
B. INSECT RESIS	TANCE: (0 = Not 1	Fested, 1 = Susceptible, 2	? = Resistant)				•		-
1 THRIPS				0 8	URROWING BU	3			
1 LEAF H	OPPER			1 "	EMATODE (Spec	ify species)			м
0 SOUTH	AN CORN ROOTW	ЮВМ		1	ESSER CORNST.	ALK BORER		1 3	
0 APHID			1		THER (Specify)	Sclero	tini RAD	<u>الط</u> حا9	<u>ight</u> 3104
. COMPARISON	OF SUBMITTED V	ARIETY WITH ONE OF	MORE SIM	LAR VA	RIETIES:			•	
VARIETY	OIL*	PROTEIN*	OLE LINOI ACID R	EIC: • LEIC LATIO	IODINE* NUMBER	SHELLING (%)	SMK**	(X) ·	MAIN STEM HEIGHT (CM)
SUBMITTED	49.5	28.04	29	.0	76.8	72.9	68.2	21	43.6
SIMILAR	47.4	27.68	28	.8	76.0	75.6	71.0	19	37.6
NAME OF SIMILA	TamrunOL01	1. 1	SunOlei			R Florun.	Floru.	Flor.	Florunner
* From Sound Mai	******	** Sound Mature Kerr		<u> </u>	ge Kernels	<u> </u>			
	VARIETY WHICH	MOST CLOSELY RESE VARIETY	MBLESTHA	I SUBMI	TTED: CHARACTER		VAS	RIETY	
OO COLOR				SEEDL	NG VIGOR		Tamru		·
EED DORMANC	,	Florunner			HICKNESS				
EEO SIZE		Florunner Tamrun 96		LEAF C	·····		Tamru Tamru		
11. COMMENTS	Additional descripti	ion or darification — Su	ch es: Relativ			compered with a			 .

Exhibit D.

ADDITIONAL DESCRIPTION OF 'TAMRUN OL 02'

Yield, Grade and Value per acre:

Tx977053 has been yield tested under non-disease conditions, TSWV infection, sclerotinia blight (Sclerotinia minor Jagger) infection and other diseases in several areas of Texas and in two research locations in Oklahoma. In 21 tests in Texas and Oklahoma, 1998 - 2001, Tx977053 averaged 10% higher pod yield (Table 1) than Florunner. These 21 tests were all three or four rep. tests with two-row plots, 10 to 15 feet long; 20 entries per test. The only major acreage of a high O/L variety being grown at this time is 'Flavor Runner 458' which is basically a high O/L Florunner that has limited disease resistance. Tx977053 has shown an average 21% yield advantage over Flavor Runner 458 in 21 tests between 1999 and 2001 (Table 1). This included a 5.7% average yield advantage in nine field tests that were disease free (non-significant, DMRT p = 0.05) and a 38% advantage in seven (7) field tests with TSWV, Sclerotinia blight, and/or other diseases (significant DMRT, p = 0.05) (see data and analyses in Appendix IV). Grades (TSMK) for Tx977053 have averaged 3% lower in non-diseased fields and nearly equal in areas of disease pressure (Table 1).

Several other comparisons are shown in Table 1 for West Texas, Oklahoma, Central, and South Texas. The yield and grade factors are about the same in all these tests. Tx977053 was in the top yielding class in most all tests, and often averaged higher than most varieties, lower in others, but only in the three-test averages at Ft. Cobb, OK was Tx977053 not in the top statistical group in yield per acre.

Disease evaluation:

Our disease evaluations of Tx977053 have been made at four locations. The TSWV evaluations have been conducted mostly in South Texas, near Pearsall, but some limited data has been collected at Stephenville, Texas. Sclerotinia evaluations were made at Stephenville and at Ft. Cobb, Oklahoma. Limited southern blight evaluations were made at Yoakum, Texas. Unfortunately, drought and confounding by other diseases at Stephenville have hampered some tests. The data are shown in **Appendix IV**.

Some statements regarding these and related data (some not shown) are appropriate, as follows:

- TSWV ratings of Tx977053 have been consistently better than the susceptible 'Tamrun 88'.
- TSWV ratings for Tx977053 have been better numerically than for Florunner in three of five tests.
- Two of three ratings for rust were better for Tx977053 than Florunner.
- -Averages for Tx977053 for Sclerotinia ratings were consistently lower (lower is better) than Florunner, Okrun, and Flavor Runner 458, however most of the differences were not statistically significant, so it is difficult to make definitive decisions based solely on these data. (See Appendix IV).

Our observations of Tx977053 indicate that the line will perform better than Florunner and Flavor Runner 458 in fields infested with TSWV, sclerotinia, southern blight and rust (Appendix IV). However, the yield differences between Tx977053 and Flavor Runner 458 under diseased conditions were significant (Appendix IV), with TX977053 having a 38% advantage.

Table A. Data for main stem height, lateral branch length and leaflet length and width from three locations.

Line	MS hth	Lat. Lgt.	Leaflet Igt.	Leaflet wdt.
	cm	cm	mm	mm
TX977053	43.6 a **	61.6 a	59.7 a	22.0 a
Florunner	37.6 b	55.4 a	45.9 c	18.9 b
Tamrun 96	31.5 c	44.9 b	48.2 c	20.0 b
Tamrun OL01	29.4 с	45.6 b	50.7 b	18.4 b

Table B. Pod and seed measurements.

Tx977056, Tamrun OL01, Florunner, & Tamrun 96 Pod and seed measurements in mm. Summary*

Tx977053	Tamrun OL01	Florunner	Tamrun 96	CV
31.4 b**	33.6 a	28.6 d	29.4 c	9.98
11.6 b	12.8 a	11.4 b	11.7 b	8.39
12.1 b	13.6 a	11.8 b	11.5 b	8.04
11.8 b	13.2 a	11.6 b	11.6 b	7.96
8.8 b	10.5 a	9.1 b	9.1 b	13.2
16.3 a	17.6 a	14.3 b	14.0 b	10.8
15.2 b	17.0 a	12.5 c	13.0 c	9.6
15.9 b	17.3 a	13.6 с	13.5 с	10.4
8.85 a	8.4 a	7.6 b	8.7 a	11.9
9.31 b	9.39 a	7.9 c	8.5 c	11.2
9.08 a	8.94 a	7.8 b	8.6 a	11
	31.4 b** 11.6 b 12.1 b 11.8 b 8.8 b 16.3 a 15.2 b 15.9 b 8.85 a 9.31 b	31.4 b** 33.6 a 11.6 b 12.8 a 12.1 b 13.6 a 11.8 b 13.2 a 8.8 b 10.5 a 16.3 a 17.6 a 15.2 b 17.0 a 15.9 b 17.3 a 8.85 a 8.4 a 9.31 b 9.39 a	31.4 b** 33.6 a 28.6 d 11.6 b 12.8 a 11.4 b 12.1 b 13.6 a 11.8 b 11.8 b 13.2 a 11.6 b 8.8 b 10.5 a 9.1 b 16.3 a 17.6 a 14.3 b 15.2 b 17.0 a 12.5 c 15.9 b 17.3 a 13.6 c 8.85 a 8.4 a 7.6 b 9.31 b 9.39 a 7.9 c	31.4 b** 33.6 a 28.6 d 29.4 c 11.6 b 12.8 a 11.4 b 11.7 b 12.1 b 13.6 a 11.8 b 11.5 b 11.8 b 13.2 a 11.6 b 11.6 b 8.8 b 10.5 a 9.1 b 9.1 b 16.3 a 17.6 a 14.3 b 14.0 b 15.2 b 17.0 a 12.5 c 13.0 c 15.9 b 17.3 a 13.6 c 13.5 c 8.85 a 8.4 a 7.6 b 8.7 a 9.31 b 9.39 a 7.9 c 8.5 c

^{*}This is a summary of 11 separate analyses conducted on data collected from three locations.

^{**}Means followed by same letter in same row are not different DMRT (0.05)

Table C. Comparative data of conventional and high O/L
Peanut breeding lines grown in South and West Texas
(Unpublished data)

Oleic/Linoleic Acid Ratios 1993 - 1995 Crop Years

Breeding Line	Frio Co. Average 1994-95	Brazos Co. Average 1993-95	Gaines Co. Average 1994-95
Tx932802 ⁴ Tx932808 ⁴		1.35	1.18
		20.26	15.56
Tx932821*		14.27	10.56
Tx932857*	,	1.25	0.99
Tx932894*		16.81	12.55
Tx932904*	1.33	1.27	1.05
Average	8.65	7.78	5.89
	Frio Co.	Brazos Co.	Gaines Co.
	% Frio Co.	% Frio Co.	% Frio Co.
Tx9328021	100.00	09.00	
Tx932808*	100.00	98.96	86.05
Tx932821 ¹		108.52	83.35
•	100.00	80.93	59.88
Tx932857*	100.00	98.31	77.62
Tx932894'	100.00	74.74	55.81
Tx932904*	100.00	95.39	79.29
	100.00	89.95	68.12

J. LEEK ASSOCIATES, INC.

LOND Probability Proces Albury, Georgia (470) CDM (83) (826)

Albany, Georgia 51705-0°529 fax: (220) 838-1166 P.O. POS. 2003; S.

Fatty Acid Composition Results 01/08/2002

	1			- O/L. Value	FAC180	FAC180	FAC181	FAC182	FAC183	FAC200	FAC201	EACOON	EAC224	EAC240
	Sepu	Unknown	Jumbo	10.8 5.31	5.31	2.23	75.58	7.02		1.32	2.52	300	7 7	2.15
	2 Steph	Florunner Medium	Medium	1.3	9.32	•	44.76	35.40	000	1.32	180	3.30	0 19	2.12
·	3 Steph	Tamrun 96 Jumbo	Jumbo	1.4	9.20	2.40	46.65	33.34	00.0	1.72	146	3.29	000	1.95
(4 Steph	Tamrun 96 Medium	Medium		9.13	2.32	45.14	34,53	00.0	1.55	1.68	3.54	0.00	2.1
Duplicat Steph	t Steph	Tamrun 96 Medium	Medium	1.3	8.96	2.30	45.12	34.34	00:0	1.63	1.70	3.73	0.00	2.23
	5 Steph	Flv.Ru.458 Jumbo	Jumbo	9.7	5.41	1.70	76.04	7.82	00.00	1.17	2.47	3.00	0.28	2.11
9	Steph	Flv.Ru.458 Medium	Medium	8.2	5.48	1.55	74.16	8.99	0.00	1.41	2.73	3.09	0.37	2.22
_	7 Steph	TR OL01	Jumbo	18.2	5.04	2.04	79.46	4.36	0.00	1.45	2.68	2.97	0.25	2.04
æ ,	8 Steph	TR OL01	Medium	6.6	5.26	1.77	75.44	7.61	0.00	1.29	2.85	3.14	0.45	2.19
ை	9 Steph	Tx977053	Jumbo	18.0	5.37	2.00	79.69	4.43	0.00	1.47	2.17	2.83	0.24	1.81
10	0 Steph	Tx977053 Medium	Medium	17.7	5.23	1.85	78.37	4 44	0.00	1.36	2.63	3.46	0.32	2.35
	1 Frio Co	Florunner	Jumbo	1.6	9.37	2.04	48.78	31,12	0.28	1.42	1.50	3.19	0.15	2.01
12	12 Frio Co	Florunner Medium	Medium	1.5	9.40	1.97	48.32	31.47	0.00	1.51	1.60	3.43	0.17	2.11
Duplicat	Duplicati Frio Co	Florunner Medium	Medium	1.6	9.42	2.02	48.36	31.06	0.00	1.39	1.67	3.54	0.17	2.15
	3 Frio Co	Tamrun 96 Jumbo	Jumbo	1.7	9.35	2.53	50.28	29.43	0.00	1.75	1.38	3.31	0.00	1.97
4	Frio Co	Tamrun 96 Medium	Medium	1.6	9.31	2,45	48.93	31.05	0.00	1.57	1.47	3.25	0.00	1.97
15	Frio Co	TR OL01	Jumbo	17.9	5.81	2.16	79.08	4.41	0.00	4.	2.04	2.90	0.21	1.95
16.	Frio Co	TR OL01	Medium	15.5	5.55	2.01	77.16	4.99	0.00	1.42	2.34	3.09	0.27	2.98
17	17 Frio Co	Tx977053	Jumbo	19.8	5.66	2.08	79.32	4.01	0.31	1.41	2.09	2.99	0.20	1.95
18	Frio Co	18 Frio Co Tx977053 Medium	Medium	16.4	5.74	1.92	78.18	4.78	0.00	1.41	2.25	3.26	0.24	2.10
19	West TX	Florunner	lumbo	4.	86.8	2.05	47.55	32.92	0.00	1.54	1.63	3.26	0.00	2.07
28	1	TX Florunner Medium	Medium	4.	9.23	2.03	46.01	34.02	0.00	1.39	1.68	3.36	0.17	2.11
Duplicat West		X Florunner Medium	Medium	1.3	9.29	2.02	45.67	34.07	0.20	1.31	1.66	3.31	0.17	2.14
21	_	-X Tamrun 96 Jumbo	oquin	4.	80.6	2.63	45.95	33.33	0.00	1.70	1.54	3.45	0.16	2.16
22	22 West TX	Tamrun 96 Medium	/edium	1 .3	9.36	2.55	44.96	33.64	0.10	1.51	1.66	3.60	0.18	2.24
23	_	X TR OL01 Jumbo	oqun	15.0	5,15	2.45	77.57	5.18	0.27	1.47	2.19	2.93	0.24	2.12
24	_	X TROLO1 IN	Medium	13.6	5.22	2.32	77.88	5.73	0.00	1.45	2.38	2.74	0.26	2.02
25	25 West TX		oquin	19.5	5.28	2.40	80.08	4.10	0.00	1.30	2.07	2.70	0.21	1.85
26 1	26 West TX	TX Tx977053 1	Modium	19.9	5.16	2.30	79.79	4.01	0.19	1.35	2.21	2.88	0.24	1.90

Table D. Percentage of Oleic and Linoleic acid in Tx977053 Samples from 2001 crop year

Average for 100 randomly chosen seeds O/L ratio = 29.3 %Oleic acid = 86.24 % Linoleic acid = 3.1

Average for 100 seeds but eleminate any under ratio of 18:1.

O/L ratio = 29.6

% Oleic acid = 86.35

% Linoleic acid = 3.0

Additional data for Tx977053 are presented in Appendix III.

Table E.

Sugar Content Tx977053 2001

Measurements in percent

	West TX ALT	South TX ALT	WTPGRF ALT	Mean
Tx977053	4.8	3.9	5.2	4.63 c*
Florunner	4.5	3.8	5	4.43 d
Tamrun OL01	5.4	4.4	5.9	5.23 a
Tamrun 96	4.9	4.1	5.3	4.77 bc
Flv. Run. 458	4.9	4.2	5.6	4.90 b

^{*}Means followed by same letter are not different, DMRT (0.05)

Table 1. Comparison data on high O/L Tamrun OL01 and Tx977053 1998 to 2001

Yield, TSMK, 100 sd. wt., and Value/Acre

	Pods/Acre	TSMK	100 Seed	Val./Acre*	Disease
Variety/					
Breeding line	lbs.	%	g	\$	%
	Nest Texas 200	 			
Florunner	4300 a**	76.1 a	64.4 b	1413	
Tx977053	4123 a	71.2 b	67.8 b	1272	
Tamrun OL01	3767 a	73.2 b	76.8 a	1193	
4 Tests - V	West Texas 200	0 - 2001			
Florunner	4447 a	78 a	66.1 bc	1466	
Flavor Runner 458	4411 a	76 a	63.3 c	1466	
TX977053	4289 a	72 b	69.9 b	1330	
Tamrun OL01	3873 a	74 ab	76.4 a	1236	
3 Tests - F	t. Cobb, Oklah	oma 2000 - 2	001		
Tamrun OL01	3830 a	69.6 a	70.7 a	1140	19.3**
Tx977053	3384 b	70.2 a	63.2 ab	1026	25.8**
Florunner	2882 с	70.1 a	55.6 bc	873	31.7**
Flavor Runner458	2779 с	71 a	53.7 c	776	39.4**
14 Tests -	All areas All	including Tx9	977053 and Fl	avor Runner	458
Tamrun OL01	4502 a	70.2 bc	71.9 a	1,370	
Tx977053	4523 a	68.8 c	64.9 b	1,349	
Tamrun 96	4438 a	68.9 c	60.5 c	1,318	
Florunner	4176 ab	71.6 ab	58.8 cd	1,299	
Flv. Runner 458	3921 b	72.3 ab	57.7 d	1,232	

^{*} Value determined by using Price Schedules for each year.

^{**} Means followed by the same letter within the column and within one test site are not significantly different. Duncan MRT 0.05.

^{**} Average percent sclerotinia diseased plants per plot

		. 7053 REL	EASE	10	:45 Mond	lay, July 1	, 2002 9	
		eneral Linear Mod	lels Procedu	ire	÷			
Dependent Variab	10: 100 sd	aut						
Source	DF	Sum of Square	es	Mean	Square	F Value	Pr > F	,
Model	22	3274.377777	78	148.8	3535354	19.91	0.0001)
Error	40	299.0663492	21	7.4	7665873			
Corrected Total	62	3573.4441269	98	,		•		
	R-Square	C.\	<i>i</i> .	, R	oot MSE	•	YIELD Mean	
	0.916309	4,14195	55	2.7	3434795	6	6.01587302	
		.*	•					
Source	DF	Type I S	SS	Mean	Square	F Value	Pr > F	
T LOC	2 20	2013.7536507 1260.6241269			7682540 3120635		0.0001)
Source	DF	Type III 8	ss .	Mean	Square	F Value	Pr > F	
roc 1	2 20	2013.753650 1260.6241269			7682540 3120635		0.0001)
		7053 REL	EASE	10:	:45 Mond	lay, July 1	. 2002 10	
	G	eneral Linear Mod	els Procedu				•	
		Multiple Range Te			IELD 00	st wit		
	NOTE: This test experimen	controls the typ twise error rate	e I comparis	sonwise	error r	ate, not t	he	
	Al	pha= 0.05 df= 40	MSE= 7.476	6659	-		`.	
٠		Number of Means Critical Range	2 3 1.705 1.793					
	Means with the	same letter are n	ot significa	antly di	ifferent			
. •	Duncan (Grouping	Mean	N T	ī			
		Α .	73.4952	21 T	ROLO1	Tamor C	9L 01	
	·	В	64.7238	21 T	ROLO2			
		С	59.8286	21 F	LO	Florum	4	

				-	,					TE VE
		7053	RELEASE	10 165 1	:45 Mond	ay, July 1	, 2002	6		
	OBS	T	LOC	100 yang	<u>'</u>					
. •	1	FLO	1	62.5						
	2	FL0	2	65.5		7053 R	ELEASE			Monday, July
	3	FLO	3	64.2				100	4	
	. 4	FLO	4	57.2	OBS	Ť	LOC	Y255C-	ant	, etc.
	5	· FLO	5	53.9		T00100		FO F		
	6	FLO	6	59.7	52	TROLO2	10	58.5		-
	7	FLO	7	69.0	53	TROLO2	11	70.2		
	8	FLO	8	72.6	54 55	TROLO2 TROLO2	12 13	63.0 65.3		
	9	FLO	9 .	58.6	56	TROLO2	14	59.5		
	10	FLO	10	51.0	57	TROLO2	15	67.9		
	11	FLO	11	59.2	58	TROLO2	16	63.1		
	12	FLO	12	56.7	59	TROLO2	17	59.1		
	13	FLO	13	62.3	60	TROLO2	18	64.5		
	14	FLO	14	62.1	61	TROLO2	19	61.0		•
	15	FLO	15	59.9	62	TROLO2	20	63.9		
	16 17	FLO	16	54.6	63	TROLO2	21	69.1		•
	18	FLO FLO	17	53.9	•					
	19	FLO	18 19	. 58.8 56.5						
	20	FLO	20	57.3						
	21	FLO	21	60.9				\		
	22	TROLO1	1	77.1						
	23	TROLO1	2	76.4						
	24	TROLOI	3	75.3	`					
	25	TROLO1	4	73.8						
·	26	TROL01	5	63.7		*				
	27	TROL01	6	77.9						
	28	TROLO1	7	79.6)
	29	TROL01	. 8	77.6						
	30	TROLO1	9	77.4						
	31	TROL01	10	61.4						
	32	TROLO1	11	79.8	-					
	33	TROL01	12	66.6						
	34	TROLO1	13	76.7						
	35	TROLO1	14	78.2						•
	36	TROLO1	15	72.6						
	37	TROLO1	16	72.4						
	38	TROLO1	17	66.5		• .				•
	39	TROLO1	18	71.0					=	
	40	TROLO1	19	71.0						
	41	TROLO1	20	68.4					•	
	42 43	TROLO1 TROLO2	21 1	80.0 61.9						· ·
	44	TROLO2	. 2	64.5					•	•
	45	TROLO2	3	65.9					•	
	46	TROLO2	4	63 1						•
	47	TROLO2	5	56.9						
	48	TROLO2	6	66.0		*				
	49	TROLO2	7	72.0				. '		•
	50	TROLO2	8	74.5						
	51	TROLO2	.9	69.3						. ,
in the second	an E value	ingings.								

100 Seed weight for Tamrun OL 02 PVP application No. 200300170

Location No.	Loc. ID	Test year
1.	Frio Co. Neal Farm	1998
2.	Frio Co. Phillips Farm	1998
3.	Frio Co. Neal Farm	1999
4.	Frio Co. Phillips Farm	1999
5.	Frio Co. Neal Farm	2000
6.	Frio Co. Phillips Farm	2000
7.	Gaines Co. Johnson Frm.	2000
8.	Gaines Co. WTPGRF	2000
9.	Erath Co W/O Disease	2000
10.	Ft. Cobb Okla. Ham	2000
11.	Ft. Cobb Okla. Keim	2000
12.	Erath Co. 2001 UPPT	2001
13	South Texas UPPT	2001
14.	West Texas UPPT	2001
15.	Frio Co. Phillips Farm	2001
16.	Frio Co. Neal Farm	2001
17.	Erath Co. W Disease	2001
18.	Erath Co. W/O Disease	2001
19.	Ft, Cobb Ham	2001
20.	Gaines Co. WTPGRF	2001
21.	Gaines Co. Johnson Farm	2001

		antiferrom to	linging lands a S					4120 00000	unikani suku
		e west			(d. 1911)		estaplement	nenga Alegiya	
and the contraction							generalisas eser		
								100	
	- 112	# 201	00 ALT 10	a en un	r 09:27 I	Eniday	Contomb	an 94 00	104 - 2
Tes+1	Evath Cones	<i>"</i> "	JO ALI 10			r tuay,	gehrenn	EI 24, 23	04 2
145+1			Γ	kep Loc	<i>≸eu</i> SDWT			All Control	
			TROLO2	1	69.5				1.3
			FROLO2	2	71.6				
			FROLO2 FR96	3	76.7				
			rneo rree	1 2	68.6 69.7				
			TR96	3	68.7	1000			
			LO	1	62.7				
		8 F	TLO	2	52.3		Maria de la composição de La composição de la compo		
		9 F	LO	3	60.7				
,		Th	ie GLM Pro	neadura					
			O MEM III	rcedul e					
Dependent Variat	ole: SDWT			-					
						4.75			
Pauma			Sum					Type 19 Type 19	
Source		DF	Squar	'es	Mean Squa	re F	Value	* Pr > F	
Model		4	044 00444					4. 1. 1.	
		4 ;	344.99111	11	86.24777	78	5.49	0.0639	: .
Error		4	62.82444	.44	15.70611	44			
		1.			10.70011				
Corrected	Total	8	407.81555	56	100				•
	n n-				a gray			5 A 4 A 4 A 5 A 5 A 5 A 5 A 5 A 5 A 5 A	
	R-Square	Coeff	Var	Root N	ISE SDI	WT Mean	4,	11 N	
	0.845949	5.939	 						
	51.4.50	0.505	 	3.9630	194 60	5.72222	4.1.		
		:							
Source		F	Type I S	38	Mean Squar	e F	Value -	Pr > F	
·						•	,44200		
toc Rep			18.748888		159.374444	14	10.15	0.0271	$\overline{}$
		2	26.242222	22	13.121111	11	0.84	0.4975	
					4 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1				et e
Source	D	FT	ype III S	100	Mean Pau-) (m.2)	_	
We would be a second		٠,	Abe iff o	,c)	Mean Squar	e F	Value	Pr > F	-
T		2 3	18.748888	9	159.374444	4	10.15	(0.0271	> -
Lookep			26.242222		13.121111		0.84	0.4975	
1	and the second s	and the second	100	M		Programme Company		J. TJ. J	医线性性皮肤

Test 1

Erath County

2000 ALT 100 SD WT 09:27 Friday, September 24, 2004

The GLM Procedure

Duncan's Multiple Range Test for SDWT

NOTE: This test controls the Type I comparisonwise error rate, not the experimentwise error

Alpha

Error Degrees of Freedom Error Mean Square

15.70611

Number of Means Critical Range

8.984

9.181

Means with the same letter are not significantly different.

	100 Seed	wt		
Duncan Grouping	Mean	'N	T	
А А	72.600	3	TROLO2	Tamum 0L02
Â	69.000	3	TR96	Tamum 96
В	58.567	3	FLO	Florumer

Test 2						
1 TROLO2	to + 1	• .				ber 24, 2004
1 TROLO2	1th 2	Obs	T Re	o /eo - SDWT	ing the second	
2						
3		of the contract of the contrac	and the second of the second o			
Source DF Sum of Square Square		3	TR0L02 3	tine that a parties, and a first the first terms of the contract of the contra		
Corrected Total 11 133.8900000 Source DF Type I S Mean Square F Value Pr > F		The same of the sa		The committee of the property of the competitions		
B TR96		6	TR96 2			
9 FLO 1 58,5 10 FLO 2 57,9 11 FLO 3 56,2 12 FLO 4 62,6	en e	2.11	医乳腺性 医电影性 化二氯甲基甲基甲基甲基甲基甲基甲基甲基甲基甲基甲基甲基甲基甲基甲基甲基甲基甲基甲基	机工程设置 化基苯酚 化二氯甲酚 化二氯甲酚二二氯甲酚		
11		A CONTRACTOR OF THE STATE OF				
The GLM Procedure Dependent Variable: SDWT Source DF Squares Mean Square F Value Pr > F Model 5 103.5650000 20.7130000 4.10 0.0577 Error 6 30.3250000 5.0541667 Corrected Total 11 133.8900000 R-Square Coeff Var Root MSE SDWT Mean 0.773508 3.725182 2.248147 60.35000 Source DF Type I SS Mean Square F Value Pr > F T 2 56.73500000 28.36750000 5.61 0.0423 3 46.83000000 15.61000000 3.09 0.1114 Source DF Type III SS Mean Square F Value Pr > F T 3 6 73500000 28.36750000 3.09 0.1114			The second secon	the state of the s		
Source DF Squares Mean Square F Value Pr > F						
Source DF Squares Mean Square F Value Pr > F						
Source DF Squares Mean Square F Value Pr > F Model 5 103.5650000 20.7130000 4.10 0.0577 Error 6 30.3250000 5.0541667 Corrected Total 11 133.8900000 R-Square Coeff Var Root MSE SDWT Mean 0.773508 3.725182 2.248147 60.35000 Source DF Type I SS Mean Square F Value Pr > F T 2 56.73500000 28.36750000 5.61 0.0423 3 46.83000000 15.61000000 3.09 0.1114 Source DF Type III SS Mean Square F Value Pr > F	•		The GLM Procedu	re		
Source DF Squares Mean Square F Value Pr > F Model 5 103.5650000 20.7130000 4.10 0.0577 Error 6 30.3250000 5.0541667 Corrected Total 11 133.8900000 R-Square Coeff Var Root MSE SDWT Mean 0.773508 3.725182 2.248147 60.35000 Source DF Type I SS Mean Square F Value Pr > F T 2 56.73500000 28.36750000 5.61 0.0423 3 46.83000000 15.61000000 3.09 0.1114 Source DF Type III SS Mean Square F Value Pr > F	Dependent Variable: SDWT					•
Source DF Squares Mean Square F Value Pr > F Model 5 103.5650000 20.7130000 4.10 0.0577 Error 6 30.3250000 5.0541667 Corrected Total 11 133.8900000 R-Square Coeff Var Root MSE SDWT Mean 0.773508 3.725182 2.248147 60.35000 Source DF Type I SS Mean Square F Value Pr > F T 2 56.73500000 28.36750000 5.61 0.0423 L00 Rep 3 46.83000000 15.61000000 3.09 0.1114 Source DF Type III SS Mean Square F Value Pr > F	Separative for Education of the		· ·			4
Error 6 30.3250000 5.0541667 Corrected Total 11 133.8900000 R-Square Coeff Var Root MSE SDWT Mean 0.773508 3.725182 2.248147 60.35000 Source DF Type I SS Mean Square F Value Pr > F T 2 56.73500000 28.36750000 5.61 0.0423 1 46.83000000 15.61000000 3.09 0.1114 Source DF Type III SS Mean Square F Value Pr > F	Source	DF		Mean Square	F Value	Pr > F
Corrected Total 11 133.8900000 R-Square Coeff Var Root MSE SDWT Mean 0.773508 3.725182 2.248147 60.35000 Source DF Type I SS Mean Square F Value Pr > F T 2 56.73500000 28.36750000 5.61 0.0423 3 46.83000000 15.61000000 3.09 0.1114 Source DF Type III SS Mean Square F Value Pr > F T 2 56.73500000 28.36750000 5.61 0.0423 0.1114	Model	5	103.5650000	20.7130000	4.10	0.0577
R-Square Coeff Var Root MSE SDWT Mean 0.773508 3.725182 2.248147 60.35000 Source DF Type I SS Mean Square F Value Pr > F T 2 56.73500000 28.36750000 5.61 0.0423 3 46.83000000 15.61000000 3.09 0.1114 Source DF Type III SS Mean Square F Value Pr > F	Error	6	30.3250000	5.0541667		
0.773508 3.725182 2.248147 60.35000 Source DF Type I SS Mean Square F Value Pr > F T 2 56.73500000 28.36750000 5.61 0.0423 3 46.83000000 15.61000000 3.09 0.1114 Source DF Type III SS Mean Square F Value Pr > F	Corrected Total	11	133.8900000			
Source DF Type I SS Mean Square F Value Pr > F T 2 56.73500000 28.36750000 5.61 0.0423	R-Squ	are Co	eff Var Roo	t MSE SDWT M	ean	
T 2 56.73500000 28.36750000 5.61 0.0423 3 46.83000000 15.61000000 3.09 0.1114 Source DF Type III SS Mean Square F Value Pr > F	0.773	508 3.	.725182 - 2.24	18147 60.35	000	
T 2 56.73500000 28.36750000 5.61 0.0423 3 46.83000000 15.61000000 3.09 0.1114 Source DF Type III SS Mean Square F Value Pr > F			•			
100 kg 3 46.83000000 15.61000000 3.09 0.1114 Source DF Type III SS Mean Square F Value Pr > F	Source	DF	Type I SS	Mean Square	F Value	Pr > F
+00 kp 3 46.83000000 15.61000000 3.09 0.1114 Source DF Type III SS Mean Square F Value Pr > F	T _	2	56.73500000	28.36750600	5 61	(0.0423)
Source DF Type III SS Mean Square F Value Pr > F	±00 Rep					
T 2 56 73500000 29 36750000 5 64 (6 2422)	<i>d</i>					
Τ 2 56.73500000 28.36750000 5.61 0.0423 LOO REP 3 46.83000000 15.61000000 3.09 0.1114	Source	DF	Type III SS	Mean Square	F Value	Pr > F
1.00 κερ 3 46.83000000 15.61000000 3.09 0.1114	T A	9	56.73500000	28 36750000	E 61	0 0429
	too Kep					And the Control of th

Test 2 Erath County

2001 ALT 100 SD WT 09:27 Friday, September 24, 2004 13

The GLM Procedure

Duncan's Multiple Range Test for SDWT

NOTE: This test controls the Type I comparisonwise error rate, not the experimentwise error

Alpha Error Degrees of Freedom Error Mean Square 5.054167

Number of Means Critical Range 4.031

Means with the same letter are not significantly different.

Duncan Group	ina	100 Seedu		
	Α	63.425	4 TROLOZ Tamun OLO	2
	B B	58.825	4 TR96 Tamun 96	2
	В	58.800	4 FLO Florumes	

. 1	ur3	Erach	Courts	200 4 UPPT 1	100 SD 1	WT	00.07	Enddov	Contember 04	. 14
1.0	25 Σ	,	J	÷		100	.08.27	rituay,	September 24,	2004
•			0bs	Τ.	Kar.	SDWT				
				er t					× 1.	
		٠.	. 1	TROLO2	1	60.8				
			2	TROL02	2	61.B		•		
			3	TROL02	3	62.4				
	4		4	TROL02	. 4	58.3				
			5	TR96	1	51.1				
			6	TR96	2	54.9	•			
			7	TR96	3	48.8		•		
		-	8	TR96	4	52.2				
			9	FLO	1	57.4				
			10	FLO	2.	54.0			•	
			11	FLO	3	52.7				
			12.	FLO -	~ :- ,4	53.8				
	·			The GLM Po						
				INO GEM FI	-oceaur	•				
Depen	dent Variable: SC	DWT		•						
				Sur	n of					
	Source		DF	,	ares	Mean S	quare	F Val	ue Pr > F	·
	Model		- 5	185.4283	3333	37.08	56667	7.	55 0.0144	•
	Error		6	29.4883	3333	4.91	47222			
	Corrected Total		11	214.9166	667					
		R-Square	Coe	eff Var	Root	MSE	SDWT	Mean		
		0.862792	3.	.981294	2.21	6917	55.6	8333		
					•			•		
	Source		DF	Type 1	SS	Mean S	quare	F Val	ue Pr > F	
	т		2	173.4716	1887	PE 72	58333	17.	65 (0.0031)	
	LOO Rep		3	11.9566			55556	0.		
			J	.,	,	, 0,00		0.	0.0020	
	Source		DF	Type III	SS	Mean S	quare	F Val	ue Pr > F	
	Т		2	173.4716		aa =-				
	1 .		•	173 4716					RE // NACTI	1
	too Rep		3	11.9566		86.73	55556	17.		

Fest 3 Grath Gung 200 UPPT 100 SD WT

09:27 Friday, September 24, 2004

The GLM Procedure

Duncan's Multiple Range Test for SDWT

NOTE: This test controls the Type I comparisonwise error rate, not the experimentwise error rate.

> Alpha 0.05 Error Degrees of Freedom 4.914722 Error Mean Square

Number of Means Critical Range 3.836 3.975

Means with the same letter are not significantly different.

Duncan Grouping	100 Seed wit	N	T	·
A	60.825	4	TROLO2	Tamum OLO2
8 8	54.475	4.	FLO	Florunner
B	51.750	4	TR96	Tamuan 96

James 0L02

Appendix I

Plant data

112

Duncan's Multiple Range Test for Mainstern Height

NOTE: This test controls the Type I comparisonwise error rate, not the experimentwise error rate.

> Alpha 0.05 Error Degrees of Freedom Error Mean Square Harmonic Mean of Cell Sizes 9.869159

NOTE: Cell sizes are not equal.

Number of Means Critical Range 3.692 3.880

Means with the same letter are not significantly different.

Duncan Grouping	Mean	N	т
A	43.636	33	70531 = TX477163
В	37.625	8	FLO
c c	31.500	8	TR96
· c	29.375	Ľ	7006 = Tammun OLOI → nean of 8 × 10 = 80 plants

7053 COMB 1&2 MAINSTEM MEASURMENT

15:01 Wednesday, November 14, 2001

The GLM Procedure

Dependent Variable: Win Main Slem

· ····································	HEDD WAN	course		•		
Source		DF	Sum of Squares	Mean Square	F Value	Pr > F
Model		23	2454.461404	106.715713	6.57	<.0001
Error		33	536.100000	16.245455		
Corrected Total	l	56	2990.561404			
	R-Square	1)	MSE YIELD Me		
Source		DF	Type I SS	Mean Square	F Value	Pr > F
T LOC		3 20	1915.175040 539.286364	638.391680 26.964318	39.30 1.66	<.0001 0.0960
Source		DF	Type III SS	Mean Square	F Value	Fr > F
T Loc		3	1035.600000	345.200000	21.25	<.0001

Lateral Bronch longth Duncan's Multiple Range Test for

NOTE: This test controls the Type I comparisonwise error rate, not the experimentwise error rate.

> Alpha Error Degrees of Freedom Error Mean Square Harmonic Mean of Cell Sizes 9.869159

NOTE: Cell sizes are not equal.

Number of Means Critical Range 9.78

Means with the same letter are not significantly different.

		-
A 61.606	33	70531 =TX 972053
A 55.375	8	FLO
B 45.625	8	7006 = Tanum 0601
B 44.875	8	TR96 -> meanoy 8×10 = 80 plants
B 45.625	8	FLO 7006 = Tamun CLOI

7053 COMB 1&2 LATERAL MEASURMENT

15:01 Wednesday, November 14, 2001

The GLM Procedure

Dependent Variable: Lateral Brush Length	toryth
--	--------

	-					
Source		DF	Sum of Squares	Mean Square	F Value	Pr > F
Model		23	4345.514693	188.935421	1.83	0.0550
Error		33	3407.362500	103.253409		
Corrected To	tal	56	7752.877193			
	R-Square	- (ff Var Root			
Source		DF	Type I SS	Mean Square	F Value	Pr > F
T		3 20	2890.373405 1455.141288	963.457802 72.757064	9.33 0.70	0.0001 0.7933
Source		DF	Type III ss	Mean Square	F Value	Pr > F
T		3 20	2413.837500 1455.141288	804.612500 72.757064	7.79	0.0005

Duncan's Multiple Range Test for Length length

NOTE: This test controls the Type I comparisonwise error rate, not the experimentwise error rate.

Alpha 0.05
Error Degrees of Freedom 163
Error Mean Square 31.36788
Harmonic Mean of Cell Sizes 45.40541

NOTE: Cell sizes are not equal.

Number of Means 2 3 4 Critical Range 2.321 2.443 2.524

Means with the same letter are not significantly different.

Duncan Grouping	Mean	N	Т
A	59.731	105	70531 = TA977053
В	50.700	40	7006 = Tamun 0L01
c c	48.225	40	TR96
C	45.914	35.	TR96 FLO -> mlang 35x10=350 legeleti

7053 COMBINATION GROUP 1&2 lEAF LENGTH 21 15:01 Wednesday, November 14, 2001

The GLM Procedure

Dependent Variable: West Light finall

	•	•			
Source	DF	Sum of Squares	Mean Square	F Value	Pr > F
Model	56	10234.53667	182.75958	5.83	<.0001
Error	163	5112.96379	31.36788		
Corrected Total	219	15347.50046	·		
R-Square 0.666854	(5.60		95	
T		Type I SS	Mean Square	F Value	Pr > F
LOC	3 53	7497.819740 2736.716929	2499.273247 51.636168	79.68 1.65	<.0001 0.0094
Source	DF	Type III SS	Mean Square	F Value	Pr > F
T LOC	3 53	6764.038609 2736.716929	2254.679536	71.88 /	<.0001

7053 COMBINATION GROUP 1&2 LEAF WIDTH)
15
15-01 Wednesday, November 14, 2001

The GLM Procedure

200300170

Duncan's Multiple Range Test for was leaflet with

NOTE: This test controls the Type I comparisonwise error rate, not the experimentwise error

Alpha 0.05 Error Degrees of Freedom 163 Error Mean Square 12.61658 Harmonic Mean of Cell Sizes 45.40541

NOTE: Cell sizes are not equal.

Number of Means Critical Range 1.549

Means with the same letter are not significantly different.

Duncan Grouping	Mean	N	T
A	22.0338	105	70531 = TX972053
В В	20.0500	40	TR96
C B	18.9143	35	FLO
, C	18.4250	40	7006 = Tamuen OL O/ mean of 40×10 = 400 Legglo

7053 COMBINATION GROUP 162 LEAF WIDTH 15:01 Wednesday, November 14, 2001

The GLM Procedure

Dependent Variable: Leaflit with

C_{i}	ι	-			Ē	
Source ·		DF	Sum of Squares	Mean Square	F Value	Pr > F
Mode1		56	1129.766679	20.174405	1.60	0.0122
Error		163	2056.502319	12.616579		
Corrected Total		219	3186.268998			
	R-Square 0.354574	Coeff 17.3	\			
Source		DF	Type I SS	Mean Square	F Value	Pr > F
T LOC		3 53	515.2580644 614.5086145	171.7526881 11.5945022	13.61 0.92	<.0001 0.6316
Source		DF	Type III SS	Mean Square	F Value	Pr > F
T LOC		3 53	470.2258612 614.5086145	156.7419537 11.5945022	12.42	<.0001

Appendix II

Pod and Seed Data

Duncan's Multiple Range Test for William

290300170

NOTE: This test controls the Type I comparisonwise error rate, not the experimentwise error rate.

Alpha 0.05
Error Degrees of Freedom 397
Error Mean Square 9.518287
Harmonic Mean of Cell Sizes 114.2857

NOTE: Cell sizes are not equal.

Number of Means 2 3 4 Critical Range .8024 .8447 .8730

Means with the same letter are not significantly different.

Duncan Grouping	Mean	N	T
А	33.6563	100	7006 = Tamuna OLOI
В	31.4200	200	7053 = Tx477053
С	29.4257	100	TR96
D	28.5850	100	FLO

7053 POD LENGTH-COMB.GR1&2

09:15 Friday, November 16, 20

The GLM Procedure

Dependent Variable: YHERD Pad Lingth

			Sum of			
Source		DF	Squares	Mean Square	F Value	Pr > F
Model	•	102	2614.113798	25.628567	2.69	<.0001
Error		397	3778.759822	9.518287		
Corrected Total		499	6392.873620			
	R-Square	Coeff		MSE YIELD Mea	an	
	0.408911	9.98	3.085	30.901	40	
Source		DF	Type I SS	Mean Square	F Value	Pr > F
T		3	1567.076538	522.358846	54.88	<.0001
LOC	·	99	1047.037260	10.576134	1.11	0.2416
Source		DF	Type III SS	Mean Square	F Value	Fr > F
T		3	1567.076538	522.358846	54.88	<.0001
LOC		99	1047.037260	10.576134	1.11	0.2416

Duncan's Multiple Range Test for

NOTE: This test controls the Type I comparisonwise error rate, not the experimentwise error rate.

Alpha 0.05
Error Degrees of Freedom 47
Error Mean Square 0.971679
Harmonic Mean of Cell Sizes 12.30769

NOTE: Cell sizes are not equal.

Number of Means 2 3 4 Critical Range .7994 .8407 .8678

Means with the same letter are not significantly different.

Duncan Grouping	Mean	N	T	
A	12.7780	10	7006 =	Tamun OLOI
B B	11.7400	10	TR96	
B B	11.5615	40	7053# =	Tx977053
В	11.4040	10	FLO	- 100 pods
		L	- 10K10	- 100 pod 2

7053 PODA WIDTH-COMB.GR1&2

09:15 Friday, November 16, 2001

The GLM Procedure

Dependent Variable: 11 dameter

		• • • •		•		
Source		DF	Sum of Squares	Mean Square	F Value	Pr > F
Model		22	39.94710229	1.81577738	1.87	0.0363
Error		47	45.66889200	0.97167855		
Corrected Total		69	85.61599429			
Source	R-Square 0.466585	(Root 07628 0.98 Type I SS	5738 11.738	29	D
T LOC		3 19	13.17768429 26.76941800	Mean Square 4.39256143 1.40891674	F Value 4.52 1.45	Pr > F 0.0073 0.1500
Source		DF	Type III SS	Mean Square	F Value	Pr > F
T LOC		3	10.29316800	3.43105600	3.53 (0.0218

Duncan's Multiple Range Test for

2003 00 17.0

NOTE: This test controls the Type I comparisonwise error rate, not the experimentwise error rate.

Alpha 0.05
Error Degrees of Freedom 47
Error Mean Square 1.455567
Harmonic Mean of Cell Sizes 12.30769

NOTE: Cell sizes are not equal.

Number of Means 2 3 4 Critical Range 0.978 1.029 1.062

Means with the same letter are not significantly different.

Duncan Grouping	Mean	N	Т
A	10.5490	10	7006 = Tamum 0601
В В	9.1270	10	FLO
B B	9.1120	10	TR96
В	8.8095	40	70531= TX977053
			40 Klo=400 pols

7053 POD CONSTRICTION-COMB. GR1&2

18

09:15 Friday, November 16, 2001

The GLM Procedure

Dependent Variable: Year Pol Construter

			Sum of			
Source		DF	Squares	Mean Square	F Value	Pr > F
Mode1		22	70.0049051	3.1820411	2.19	0.0123
Error	•	47	68.4116720	1.4555675		
Corrected Total		69	138.4165771	•		
•						
	R-Square	Coeff	Var Root	MSE YIELD Me.	an	
	0.505755	13.1	1.206	5469 9. 14 65	71	
Source		DF	Type I ss	Mean Square	F Value	Pr > F
T LOC		3	24.22852714	8.07617571	5.55	0.0024
БОС		19	45.77637800	2.40928305	1.66	0.0813
Source		DF	Type III ss	Mean Square	F Value	Pr > F
T	•	3	18.50878800	6.16959600	4.24	0.0099
LOC		1.9	45.77637900	2.40929205.		0.0055

Duncan's Multiple Range Test for WHELD

NOTE: This test controls the Type I comparisonwise error rate, not the experimentwise error rate.

Alpha 0.05
Error Degrees of Freedom 117
Error Mean Square 2.594795
Harmonic Mean of Cell Sizes 24.61538

NOTE: Cell sizes are not equal.

Number of Means 2 3 4 Critical Range .9093 .9570 .9887

Means with the same letter are not significantly different.

Duncan Grouping	Mean	N	т
A	17.3490	20	7006 = Tamun OLOI
В	15.9820	80	7053#= Tx477053
·	13.5625	20	FLO
Ċ	13.5375	20	TR96 20 X 10 = 200 seels

7053 SEEDA&B LENGTH-COMB.GR1&2

50

09:15 Friday, November 16, 2001

The GLM Procedure

Dependent Variable: West

	<i></i>	rengen					
Source	•	DF	Sum Squai		an Square	F Value	Pr > F
Model		22	384.09020	003 1	7.4586455	6.73	<.0001
Error		117	303.59097	740	2.5947946		
Corrected Total		139	687.68117	743	·		
	R-Square 0.558529	Coeff 10.40)	Root MSE 1.610837	YIELD Mean		
Source		DF	Туре І	ss Me	an Square	F Value	Pr > F
T LOC		3 19	239.02496 145.06523		9.6749881 7.6350124	30.71	<.0001 0.0002
Source		DF	Type III	SS Me	an Square	F Value	Pr > F
T LOC		3 19	281.01941 145.06523		3.6731387 7.6350124	36.10 2.94	<.0001 0.0002

200300170

THE ODN FLOCEGUE

Duncan's Multiple Range Test for seed width Chame

NOTE: This test controls the Type I comparisonwise error rate, not the experimentwise error rate.

Alpha 0.05
Error Degrees of Freedom 117
Error Mean Square 0.945445
Harmonic Mean of Cell Sizes 24.61538

NOTE: Cell sizes are not equal.

Number of Means 2 3 4 Critical Range .5489 .5777 .5968

Means with the same letter are not significantly different.

Duncan Grouping	Mean	N	T	
A	9.0884	80	7053 =	Tx977053
A A	8.9450	20	7006 =	Tamrum OLOI
A A	8.5880	20	TR96	·
В	7.8170	20	FLO	_
		L		20 × 10 = 200 red

7053 SEEDA&B WIDTH-COMB.GR1&2

66 09:15 Friday, November 16, 2001

The GLM Procedure

ependent Variable: Seed with (dumete)

idelie variable.	-		"Commente	Y		
Source		DF	Sum of Squares	Mean Square	F Value	Pr > F
Model		22	68.7834771	3.1265217	3.31	<.0001
Error		117	110.6170165	0.9454446		
Corrected Total		139	179.4004936			
	R-Square	ſ	f Var Root			
	0.383407	11.0	0.97	2340 8.8147	. 20	
Source		DF	Type I SS	Mean Square	F Value	Pr > F
T LOC		3 19	27.26736482 41.51611225	9.08912161 2.18505854	9.61 2.31	<.0001 0.0034
Source		DF	Type III SS	Mean Square	F Value	Pr' > F
T LOC		3 19	30.76801350 41.51611225	10.25600450 2.18505854	10.85 2.31	<.0001 0.0034

75

Appendix III

Chemical,
Organoleptic, and
Shelling Data

O/L Ratio of Tx977053 individual seed used for Breeder seed increase

	All samples	included		Exclude sar	nples less	than O/L 18:1
Sample #	O/L	% Oleic	% linoleic	O/L	%Oleic	% linoteic
162		83.9	3.7	22.4	83.9	3.7
163		86.1	3.9	22.2	86.1	3.9
164		83.4	4.8			
165		85.2	3.3	25.6	85.2	3.3
166		86.9	3.1	28.3	86.9	3.1
167	40.5	87.1	2.1	40.5	87.1	2.1
168	35.2	87.7	2.5	35.2	87.7	2.5
169	34	87.3	2.6	34	87.3	2.6
170	35.8	86.2	2.4	35.8	86.2	2.4
171	19.9	85.1	4.3	19.9	85.1	4.3
172	15.7	81.5	5.2			
173	22.2	85.1	3.8	22.2	85.1	3.8
174	28.5	87.3	3.1	28.5	87.3	3.1
175	31.1	85.7	2.8	31.1	85.7	2.8
176	32.6	86.3	2.6	32.6	86.3	2.6
177	22.5	85.5	6.8	22.5	85.5	2.0 6.8
178	34.6	87.1	2.5	34.6	87.1	2.5
179	27.2	85.4	3.1	27.2	85.4	3.1
180	39	88.1	2.3	39	88.1	
181	43.2	86.3	2.3	43.2	86.3	2.3
182	30.8	86.6	2.8	30.8		2
183	28.9	87.3	3		86.6	2.8
184	22.8	84.9	3.7	28.9 22.8	87.3	3
185	23.4	84.9	3.6	23.4	84.9	3.7
186	31.3	86.6	2.8		84.9	3.6
187	35.3	87.1	2.5	31.3	86.6	2.8
188	27.1	85.7		35.3	87.1	2.5
189	24	87.2	3.2 3.6	27.1	85.7	3.2
190	37.1	86.5		24	87.2	3.6
191	35.5	87.5	2.3	37.1	86.5	2.3
192	34.3	87.2	2.5	35.5	87.5	2.5
193	27.9	86.3	2.5	34.3	87.2	2.5
194	30.1	86.3	3.1	27.9	86.3	3.1
195	26.4		2.9	30.1	86.3	2.9
196		86.5	3.3	26.4	86.5	3.3
197	26.7 27	88.1	3.3	26.7	88.1	3.3
198		86.1	3.2	27	86.1	3.2
199	. 24.9 21	86.2	3.5	24.9	86.2	3.5
200		84.9	4	21	84.9	4
	24	86.9	3.6	24	86.9	3.6
201	24.8	86.7	3.5	24.8	86.7	3.5
202	33.1	87.5	2.6	33.1	87.5	2.6
203	45.2	87.7	1.9	45.2	87.7	1.9
204	24.5	86.2	3.5	24.5	86.2	3.5
205	35.4	87.3	2.5	35.4	87.3	2.5
206	24.2	85.8	3.5	24.2	85.8	3.5
207	26.6	84.9	3.2	26.6	84.9	3.2
208	25.3	86.1	3.4	25.3	86.1	3.4
209	33.6	86.9	2.6	33.6	86.9	2.6
210	29.2	85.6	2.9	29.2	85.6	2.9
				28		

Sar	mple #	O/L	% Oleic	% linoleic	O/L	%Oleic	% linoleic	
	211	24.6		3.5	24.6	85.1	3.5	
	212	22.9		3.7	22.9	83.9	3.5 3.7	
	213	27.6		3.1	27.6	85.8	3.1	
	214	22.6		3.8	22.6	85.1	3.8	
	215	23.8		3.6	23.8	85.2	3.6	
	216	30.1		2.9	30.1	86.1	2.9	
	217	30.4		2.8	30.4	86.6	2.8	
	218	33.4		2.6	33.4	87.5		
	219	30.2		2.9	30.2	86.9	2.6	
	220	21.5		4	21.5	85.6	2.9 4	
	221	24.5		3.5	24.5	85.1	3.5	
	222	29.6		2.9	29.6	86.9		
	223	32		2.7	32	87.4	2.9	
	224	24.3		3.5	24.3	85.2	2.7	
	225	25.4		3.4	2 4 .3 25.4		3.5	
	226	17.3		4.9	23.4	85.9	3.4	
	227	31.4	86.1	2.7	24.4	06.4	0.7	
	228	30	86.8	2.9	31.4 30	86.1	2.7	
	229	39	86.9	2.2		86.8	2.9	
	230	37.1	87.4	2.4	39	86.9	2.2	
	231	29.2	87.6	3	37.1	87.4	2.4	
	232	39.3	86.9	2 .2	29.2	87.6	3	
	233	23.4	86.6	3.7	39.3	86.9	2.2	
	234	25	85.6	3.4	23.4	86.6	3.7	
	235	39.7	87	2.2	25	85.6	3.4	
	236	35.3	87.1	2.5	39.7	87	2.2	
	237	42.4	85.5	2.5	35.3	87.1	2.5	
	238	35.3	87.8	2.5	42.4	85.5	2	
	239	18.3	82.3	4.5	35.3	87.8	2.5	
	240	33.7	86.9	4.5 2.6	18.3	82.3	4.5	
	241	31.8	88.1	2.8	33.7	86.9	2.6	
	242	25.3	86.3	3.4	31.8	88.1	2.8	
	243	28.1	87.1	3.1	25.3	86.3	3.4	
	244	28.5	86.7	3.1	28.1	87.1 90.7	3.1	
	245	27	86.5	3.2	28.5 27	86.7	3	
	246	35.1	87	2.5		86.5	3.2	
	247	27	85.9	3.2	35.1	87 85.0	2.5	
	248	20.3	85.5	4.2	27	85 .9	3.2	
	249	25.1	86.5	3.4	20.3 25.1	85.5	4.2	
	250	31	86.4	2.8		86.5	3.4	
	251	30.9	86.5	2.8	31 30.9	86.4	2.8	
	252	29.6	86.7	2.9	29.6	86.5	2.8	
	253	33.4	86.7	2.6	33.4	86.7	2.9	
	254	26.4	86	3.2		86.7	2.6	
	255	44.4	88.4	1.9	26.4	86	3.2	
	256	28.5	86	3	44.4	88.4	1.9	
	257	2 6 :5	86.1	3.2	28.5	86	3	
	258	31.5	86	3.2 2.7	26.5	86.1	3.2	
	259	38.2	85 <i>.</i> 9	2.7	31.5	86	2.7	
	260	32.5	86	2.6	38.2	85.9	2.2	
	261	30.3	86.6	2.8	32.5	86	2.6	
Total		2928.8	8624.2	2.6 310	30.3	86.6	2.8	
mean		29.3	86.24	3.10 3.1	2878.6	8375.5	295.1	
			30, <u>2</u> 7	J. I	29.6	86.35	3	



J. LEEK ASSOCIATES, INC. ~BY WORKING TOGETHER, IT CAN BE DONE~

PEANUT SEED PERFORMANCE GENERAL GUIDELINES

OIL QUALITY PARAMETERS

Peroxide Value (PV) is defined as the number of milli-equivalents as peroxide in 1000 grams of sample. It is an indicator of the degree of rancidity of a fat, but is not an "absolute index of rancidity". Peroxides form in direct proportion to oxygen absorption, but peroxides also break down readily. Therefore, the amount present in the fat at the time of the test is frequently much lower than the total amount formed.

Guidelines:

Raw peanuts (fresh): typically 0.4 max

PV>1.0 - need to find out what's going on

PV>2.0 - not good

PV>3.0 – almost impossible

Blanched peanuts: typically 1.0-3.0 (usually less than 2.0)

Roasted peanuts: can vary between 3 and 100, depending on how old (should be

3-5 on very fresh nuts.

Percent Free Fatty Acids (FFA) of a fat is a measure of the extent of hydrolysis (decomposition due to reaction with water) that has occurred in a fat. Grade factors decrease and free fatty acids increase in peanuts improperly warehoused and/or cured. Levels of deterioration are related to seed condition, length of storage, excess moisture, high temperature, and chemical damage. FFA is a good partner to PV for measuring fat quality deterioration.

Guidelines:

FFA < 0.03% is normal

Typically the upper limit for well-handled, fresh peanuts is 0.05%

Oleic/Linoleic Acid Ratio (O/L) is used as an indicator of oil stability and shelf life. It has been suggested that the selection of raw peanuts with low levels of linoleic acid is a means of extending product shelf life. Lower levels of polyunsaturated fatty acids provide less substrate for the enzyme lipoxygenase to catalyze oxygen. The principal substrates for lipoxygenase in plant systems are linoleic and linolinic acids. The primary products of the lipoxygenase-fatty acid reaction are lipid hydroperoxides, which may undergo further chemical oxidation or decomposition to produce a variety of secondary products such as aldehydes, alcohols, ketones, and acids. These are potential sources of off-flavors in many foods including peanuts and peanut products. Consequently, peanuts high in oleic acid and low in linoleic acid content are preferred to achieve better oil stability.

Guidelines for Non-High O/L Varieties

SE Runners: Typically 1.8-2.2 SW Runners: Typically 1.2-1.8

Virginias: Typically 1.6-2.5 heavily variety dependent

Spanish: Typically 1.1-1.5



J. LEEK ASSOCIATES, INC. ~BY WORKING TOGETHER, IT CAN BE DONE~

BLANCHABILITY AND ROAST VARIABILITY

Blanchability is very dependent on the use (i.e. salted nuts, candy, butter, etc...). Blanching is the removal of the testa or seed coat from shelled raw or roasted kernels. For raw peanuts this is usually done by "white roasting" or partially drying the peanuts and then processing the kernels through various types of blanchers which subject the kernels to thorough, but gentle, rubbing type friction action, followed by aspiration. Removal of the seed coat typically produces a weight loss of 3.0 to 3.5% for the shelled peanut kernels depending upon the peanut variety and grade being processed. There are four general blanching methods: 1) spin, 2) buff, 3) split, and 4) water. Each blanching process has unique utilization applications. Spin blanching will produce a premium cocktail peanut, while buff blanching is used more in preparing peanuts for confectionary coatings. Split blanching is used primarily for peanuts going into peanut butter or to prepare peanuts for removal of damaged kernels and foreign material by extensive electronic sorting. Water blanching is used to produce a "unique", crunchy texture required in many specialty products. Typical specifications for spin and buff blanched peanuts are as follows.

Physical Specifications		Chemical Specifications	
Whole Kernels	80-85%	Moisture	4.5-5.0%
Split Kernels	15-20%	Oil	49-51%
Rednose ¹	3.0%	Iodine Value	95
Unblanched ²	1.0%	Free Fatty Acids	0.30%
Damaged Kernels	1.0%	Peroxide Value	5.0 meg
	s, largest dian	Aflatoxin	5.0 ppb

Roast Variability is dependent upon many things (i.e., sugar, fat, and moisture distributions) that can be difficult to quantify. Roast variability is an important quality characteristic for all manufacturers when blending peanuts for their product, however, for manufacturers that produce "whole nut snacks" this becomes even more important. Peanuts used in non-coated, "whole nut snacks" must be very pleasing to the consumer's eye. Ranges in roast variability can vary among manufacturers. Our experience has shown that roast variability can be divided into three categories: 1) good, 2) fair, and 3) poor with the percentage of over roasted kernels ranging from ≤10%, 11-20%, and >20%, respectively.


SENSORY ANALYSIS


Flavor can be one of the most important characteristics of a peanut. The foods/snacks that consumers purchase that contain peanuts, just like any other food purchase, are based on repeat business. This means that consumers will purchase something that they like as long as they get what they want. Consumers of peanut products want a product that has a robust peanut flavor, not too sweet, with no offensive flavors. Other


Flavor Intensities and Attributes


Г	7	_	Т	_	-	T	_		_	Т	_	-	_	т	_			7	T.	_	·			_			
	<u>ه</u> د	2	3 0		, 2	٦		0	0	0	0	C	1.2	1.2	0	0	0	0	0	0	1.4	0	0	0	0	0	C)
	METALLIC	٥	0	٥			0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	- 1 EK	1 2	10) -	0	0	0	0	0	0	0	0	0	1	0	0	0			0	0	0	0	0	0	0	2
8	۲ د	P	c		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
90,5	<u></u>	P	0	9.0	0	2,4	1.2	1.8	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
8	} <	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
SADTUN	C	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Cac	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
WHS	3.2	3	2.8	2.8	3.2	3.2	3	3.2	2.2	3.2	2.8	3.2	3.2	2.8	3.2	3	3.4	2.4	2.2	5.6	3	2.8	3.2	2.6	3.2	3.2	4
č	2.8	3.6		1.4	5.6	2.8	1.4	3.2	5.6	1.4	-	2	1.6	2.6	2.2	7	3.4	3	2.2	2.4	2.2	1.6	3	2.6	2.2	2	9
RBG	1.2	9.0	1.6	1.8	1.4	1.2	1.6		2.6	1.4	2.2	1.4	1.8	1.4	1.4	1.2	0.8	1.6	1.6	1.2	1.8	1		0	0	1.2	2
TS.	2.4	2.4	2.8	3	2.8	3	3	3.4	~	2.8	3.2	3	<u>س</u>	3	3.2	3.2	2.2	3	3.4	3.2	2.8	4	3.2	3.2	3.2	3	3
æ	2.2	2.2	2.4	2.6	5.6	2,6	2	7.8	5.6	7.7	2.4	2.6	2.8	2.4	5.6	2.8	5.6	<u>س</u>	2.6	3	2.8	3.2	2.8	2.4	2.2	2.6	3
윤	5.4	5.6	5.4	5	5,4	5	5	5.2	2	4.8	2	5.6	2	5.4	5.2	5.2	5.4	5.4	5.4	5.2	5.2	2	5.8	5.2	5.4	5.4	9
Control	Unknown Jumbo	Florunner Medium	Tamrun 9 Jumbo	Tamrun 9 Medium	Flv.Ru.45 Jumbo	Flv.Ru.45 Medium	TR OL01 Jumbo	TR OL01 Medium	1x977053 Jumbo	1 x977053 Medium	Florunner Jumbo	Florunner Medium	Tamrun 9 Jumbo	Tamrun 9 Medium	TR OL01 Jumbo	TR OL01 Medium	Tx977053 Jumbo	Tx977053 Medium	Florunner Jumbo	Florunner Medium	Tamrun 9 Jumbo	Tamrun 9 Medium	TR OL01 Jumbo	TR OL01 Medium	Tx977053 Jumbo	Tx977053 Medium	trol
Š	1 Steph	2 Steph	3 Steph	4 Steph		6 Steph	7 Steph	8 Steph	9 Steph	10 Stepn	11 Fro Co	12 Frio Co	13 Frio Co	Frio	15 Frio Co	16 Frio Co	17 Frio Co	18 Frio Co	19 West TX	20 West TX	21 West TX	22 West TX	23 West TX	24 West TX		26 West TX	Contro


<u>ن</u>ر













S





Location

The GLM Procedure

Duncan's Multiple Range Test for Yann

NOTE: This test controls the Type I comparisonwise error rate, not the experimentwise error

Alpha 0.05 Error Degrees of Freedom 32 Error Mean Square 31066112 Harmonic Mean of Cell Sizes 10.76923

NOTE: Cell sizes are not equal.

Number of Means Critical Range 4893 5142 5305

Means with the same letter are not significantly different.

Duncan Group	ing	Mean	N	T
	A A	44891	6	TR96
B B	A	40107	15	TROL01
В	C C	37667	14	FLO
	C	34563	15	7053

7053 SUGAR

18:34 Sunday, July 28, 2002

Analysis for data in Table E.

The GLM Procedure

Duncan's Multiple Range Test for YIELD

NOTE: This test controls the Type I comparisonwise error rate, not the experimentwise error

Alpha 0.05 Error Degrees of Freedom Error Mean Square 816666.7

Number of Means Critical Range 1702 1773 1813 1837


Means with the same letter are not significantly different.

Duncan	Group	oing	Mean	N	T ,
		A	52333.3	3	TROL01
		B B	49000.0	. 3	FLVRU
	C C	В	47666.7	3	TR96
· .	Ċ		46333.3	3	7053
٠		D	44333.3	3	FLO

LOCATION 61 UPPT 2001 POD BULK DENSITY TABLE

FLORUNNER	GA942516
BULK Kg cm Kg/m ³ DENSITY 7.48 30.5 264.02	8ULK <u>Kg cm Kg/m³</u> DENSITY 6.80 30.5 240.03
DENSITY 7.48 30.5 264.02	DENSITY 6.80 30.5 240.03
GA962533	GA962569
BULK <u>Kg cm Kg/m²</u> DENSITY 6.82 30.5 240.67	BULK Ke cm Ka/m³
DENSITY 6.82 30.5 240.67	BULK <u>Kg cm Kg/m²</u> DENSITY 7.33 30.0 262.95
TP301-1-8	TX977006
BULK Kg cm Kg/m ³ DENSITY 7.20 29.1 266.64	BULK Kg cm Kg/m ³ DENSITY 6.82 30.5 240.96
DENSITY 7.20 29.1 266.64	DENSITY 6.82 30.5 240.96
TX977053	UF98326
BULK Kg cm Kg/m³ DENSITY 7.36 30.5 260.05	BULK <u>Kg cm Kg/m³</u> DENSITY 7.25 29.0 269.40
DENSITY 7.36 30.5 260.05	DENSITY 7.25 29.0 269.40
<u>UF98511</u>	UF97102
BULK <u>Kg am Kg/m²</u> DENSITY 7.30 30.0 262.04	BULK <u>Kg cm Kg/m³</u> DENSiTY 7.01 30.5 247.72
DENSITY 7.30 30.0 262.04	DENSITY 7.01 30.5 247.72
NC7	GA GREEN
BULK Kg cm Kg/m ³ DENSITY 6.46 30.5 227.98	BULK <u>Kg cm Kg/m³</u> DENSITY 7.39 30.5 261.00
DENSITY 6.46 30.5 227.98	DENSITY 7.39 30.5 261.00
TAMRUN 96	FLAYOR RUNNER 458
BULK Kg cm Kg/m³ DENSITY 7.27 29.0 269.77	BULK Kg cm Kg/m ³ DENSITY 7.28 30.0 261.13
DENSITY 7.27 29.0 269.77	DENSITY 7.28 30.0 261.13

LOCATION 61 POD BULK DENSITY



LOCATION 62 UPPT 2001 POD BULK DENSITY TABLE

FLORUNNER

BULK Kg cm Kg/m³
DENSITY 6.90 24.7 300.64

GA962533

BULK Kg cm Kg/m³ 6.90 26.6 265.74

TP301-1-8

BULK Kg cm Kg/m³ DENSITY 6.87 25.0 295.97

TX977053

BULK <u>Kg cm Kg/m³</u> DENSITY 6.84 24.4 301.57

UF98511

BULK Kg cm Kg/m³ DENSITY 6.86 23.5 314.18

NC7

BULK Kg cm Kg/m³
DENSITY 6.93 27.5 270.97

GA942516

BULK Kg cm Kg/m³
DENSITY 6.83 24.5 300.60

GA962569

 BULK
 Kg
 cm
 Kg/m³

 DENSITY
 6.90
 25.7
 289.01

TX977006

BULK Kg cm Kg/m³ DENSITY 6.89 24.5 302.82

UF98326

BULK Kg cm Kg/m³
DENSITY 6.90 23.1 321.72


UF97102

BULK Kg cm Kg/m³ DENSITY 6.88 25.3 292.57

TAMRUN 96

BULK Kg cm Kg·m³ DENSITY 6 88 23.2 319 23

LOCATION 62 POD BULK DENSITY



or the M

116.1

LOCATION 63 UPPT 2001 POD BULK DENSITY TABLE

FLORUNNER

BULK Kg cm Kg/m³
DENSITY 7.22 28.5 273.20

GA962533

BULK Kg cm Kg/m³
7.03 30.5 248.17

TP301-1-8

BULK Kg cm Kg/m³ 7.13 26.7 287.15

TX977053

BULK Kg cm Kg/m³ DENSITY 7.14 28.9 266.04

UF98511

BULK Kg cm Kg/m³ 7.22 28.8 270.39

NC7

BULK Kg zm Kg/m³ DENSITY 6.58 30.5 232.49

GA942516

BULK Kg cm Kg/m³ DENSITY 5.17 28 2 197.16

GA962569

BULK Kg cm Kg/m³ DENSITY 7.21 28.8 269.56

TX977006

BULK Kg cm Kg/m³ DENSITY 7.29 29.5 266.08

UF98326

BULK <u>Kg cm Kg/m³</u> DENSITY 6.92 27.2 274.18


UF97102

BULK <u>Kg</u> cm Kg/m³ DENSITY 6.72 28.0 257.92

TAMRUN 96

BULK Kg cm Kg/m³ DENSITY 7.10 27.5 277.77

LOCATION 63 POD BULK DENSITY



Appendix IV

Yield, Grade, Value/Acre, and Disease Evaluation Data The following pages in Appendix IV are examples of the more than 70 different analyses we conducted on Yield, grade and value per acre data from four years of testing on **Tx977053**.

Similar types of analyses were conducted on the disease evaluation data collected in four years testing on **Tx977053**.

Gereral Linear Models Procedure

Duncan's Multiple Range Test for variable:(<IELD

NOTE: This test controls the type I comparisonwise error rate, not the experimentwise error rate

Alpha= 0.05 df= 40 MSE= 180907.3

Number of Means 2 Critical Range 265.3 278.9

Means with the same letter are not significantly sifferent.

Duncan Grouping	Mean	N	Т	
A A	4767.0	21	TROLO1	
Ä	4738.5	21	TROLO2	
В	4314.4	21	FLO	

7053 RELEASE

11:43 Tuesday, July 2, 2002

Gereral Linear Models Procedure

Sum of Squares

Decendent Variable: YIELD

DF

Scunce

	DF	Sum of Squares	Mear Square	F Value	Pr >
Model	22	111830020.66666600	5083182.75757575	28.10	0.000
Enton	40	7236293.333333360	180907.33333334		
Octrected Total	62	119066314.00000000			
	R-Square	c.v.	Foot MSE		YIELD Mea
	0.939225	9.232967	425.33202717	460	6.666666
Sounce	· DF	Type I SS	Mear Squane	F Value	P r > 5
T _00	2 20	2698740.66666666 109131279.99999900	1349370.33333333 5456564.13000000	7.46 30.16	0.0018
Source	DF	Type:III SS	Mear Square	F Value	Pr > F
	2 20	2698740.66666666 109131279.99999900	1349370.33333333 5456564.33000000	7.46 30.16	0.0018
· ·	4	· · · · · · · · · · · · · · · · · · ·			\ \

10:53 Tuesday, July 2, 2002

General Linear Models Procedure

Duncan's Multiple Range Test for variable:

NOTE: This test controls the type I comparisonwise error rate, not the experimentwise error rate

Alpha= 0.05 df= 40 MSE= 3.811238

Number of Means Critical Range 1.218 1.280

Means with the same letter are not significantly different.

Duncan Grouping	Mean	N	т	
A A	71.6810	21	FLO	
Ā	70.9286	21	TROLO1	
В	69.4048	21	TROLO2	

7053 RELEASE

10:53 Tuesday, July 2, 2002

39.80076190

General Linear Models Procedure

∴∂Dependent Var	iable: WEEF 75M	H			
Source	DF	Sum of Squares	. Mean Square	F Value	<u></u> የr > F
Model	22	852.49904762	38.74995671	10.17	2.0001
Error	40	152.44952381	3.81123810		
Corrected Tota	al 62	1004.94857143			
	R-Square	C.V.	Root MSE		YIELD Mean
	0.848301	2.762417	1.95223925	7	0.67142857
Source	DF	Type I SS	Mean Square	f Value	Pr > F
Т	2	56.48380952	28.24190476	7.41	0.0040
LOC	20	796.01523810	39.80076190	10.44	0.0018 0.0001
Source	. DF	Type III SS	Mean Square	F Value	Pr > F
T	2	56.48380952	28.24190476	7.41	0.0018

796.01523810

LOC

20

0.0018

10.44

Gereral Linear Models Procedure

200300170

Duncan's Multiple Range Test for variable:

100 sout

NOTE: This test controls the type I comparisonwise error rate, not the experimentwise error rate

Alpha= 0.05 df= 40 MSE= 7.476659

Number of Means 2 3 Critical Range 1.705 1.793

Means with the same letter are not significantly different.

Duncan Grouping	Mean	V N	т
A	73.4952	21	TROLO1
В	64.7238	21	TROLO2
С	59.8286	21	FLO

7053 RELEASE

10:56 Tuesday, July 2, 2002

1006.87682540

63.03120635

134.67

8.43

0.000-

Gereral Linear Models Procedure

. ⊍ependent Var	riable: 100 /	2d wtg			İ
Source	DF	Sum of Squares	Mean Square	F Value	Pr >
Model***	22	3274.3777778	148.83535354	19.91	0.001
Error	40	299.06634921	7.47665873		
Connected Tot	al 62	3573.44412698		٠.	
	R-Square	c.v.	Root MSE	,	YIELD Vear
	0.916309	4.141955	2.73434795		6.015873C2
Sounce	DF	Type I SS	Mean Square	F Value	Pr > =
T	2	2013.75365079	1006.37682540	134.67	0.000
100	20	1260.62412698	63.03120635	8.43	0.000jr 0.000r
Sounce	DF	Type III ss	Mean Square	F Value	Pr > =

2013.75365079

1260.62412698

2

20

_00

7053 RELEASE

10:49 Tuesday, Jul; 2, 2002

General Linear Models Procedure

Duncan's Multiple Range Test for variable WD

NOTE: This test controls the type I comparisonwise error rate, not the experimentwise error rate

Alpha= 0.05 df= 40 MSE= 20245.44

Number of Means Critical Range 88.75 93.31

Means with the same letter are not significantly different.

Duncan Group	ing	Mean	.N	T
	A A	1459.71	21	TROLO1
8 8	Â	1425.14	21	TROLO2
8		1343.57	21	FLO

7053 RELEASE

10:49 Tuesday, July 2, 2002

General Linear Models Procedure

ependent Variabl	le: Hal	lac I			
Source	DF	Sum of Squares	Mean Square	F Val.e	Pr > F
odel	22	11718058.09523800	532639.00432900	26.3*	0.0001
Error	40	809817.61904763	20245.44047619		
Corrected Total	62	12527875.71428570			
	R-Square	C.V.	Root MSE	Y	IELD Mean
	0.935359	10.09499	142.28647327	1409	.47619048
Source	DF	Type I SS	Mean Square	F Value	Pr > F
ī	2	149367.71428571	74683.85714286	3.69	0.0339
LOC	20	11568690.38095230	578434.51904762	28.57	0.0001

Type III SS

149367.71428571

11568690.38095230

DF

2

20

Source

50C

Pr > F

0.0339

F Value

3.59

28.57

Mean Square

74683.85714286

578434.51904762

1998	}	•	•	:	
	South Texas	Disease Index			
		mean			
	Tamrun 96	5.0 ns			
	Florunner	4.0			
	Tamrun OL01	5.0			
	Tx977053	5.7			
1999					
	South Texas				
	<u>Neal</u>	TSWV	Rust	Southern	
		index	index	blight	
	Tx977053	4.0 ns	6.0 ns	ongin	
	Tamrun OL01	5.0	5.3		
	Tamrun 96	4.3	5.8		
	Florunner	3.5	5.8	.,	
			2.0		
	<u>Phillips</u>				•
	Florunner	3.0 ns	5.3 ns	8.8 ns	
	Tx977053	2.0	5.2	3.7	
	Tamrun OL01	2.8	5.0	10.3	
	Tamrun 96	1.8	4.8	5.1	
				J.1	
2000		TSWV			
	<u>Phillips</u>	% infection	t		
	Florunner	10.8 b*			
	Tx977053	4.9 b			
	Tamrun OL01	2.9 Ь			
	Tamrun 96	· ···- 0 a	\$		
	Flav. Run. 458	2.9 Ь			
2001	•				
2001					
	n Texas	TSWV			
Pr	nillips	index	Rust		
•	Tamrun OL01	5.25 a*	5.75 ns		
	Tx977053	4.75 a	5.0		
	Tamrun 96	4.75 a	5.5		
	Florunner	6.25 ab	5.25		
	Elv. D. 150	/ ·			

^{*}Means in the same column followed by the same letter are not significantly different, DMRT (0.05).

6.75 b

Flv. Ru. 458

5.25

Disease ratings for Tx977053 2000 - 2001 SCLEROTINIA

2000		OK ALT	Steph. ALT
		% inf. plts	index (1 to 10)
	Tx977006	13 ns	8.7 ab*
	Florunner	20	10 Ь
	Flav. Runner	23	9.3 b
	Tx977053	7 .	9.0 ab
	Tamrun 96	7	8.3 ab
	GA. Green	8	9.3 Б
	Tamrun 98	9	6.7 a

2001

Oklahoma

	Test 1	Test 2	Test 3	Test 4
	T-10.	% infected	plants	
Tamrun OL01 Tx977053 Tamrun 96 Florunner or Okrun	19.0 ? 30.0 15.0 36.5	18.2 ns 20.0 14.8 23.4	25.8 ab* 40.4 c 18.2 a 33.6 bc	33.0 ab* 42.8 bc 27.8 a 51.8 c
Flv. Run. 458	49.5		45.6 c	

Stephenville

	Sclerotinia	
	Disease index	
Tamrun OL01	5.75 ns	
Tx977053	4.75	
Tamrun 96	4.25	
Florunner	4.50	
Flav. Ru. 458	6.00	

[?] Statistics not available, only means.

Overall comparison of Tx977053 to Flavor Runner 458 for yield under disease pressure and Without disease pressure.

Seven tests with <u>Disease pressure</u>		Nine tests without Disease pressure
Tx977053	4784 a*	3985 a
Flavor Runner 458	3476 b	3769 a

^{*} Means within columns with the same letter are not significantly different, DMRT (0.05).

^{*}Means in the same column followed by the same letter are not significantly different, DMRT (0.05).

REPRODUCE LOCALLY. Include form number and edition date on a	il repraductions. F	ORM APPROVED - OMB No. 0581-005
U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE Application is required in order to determine if a p certificate is to be issued (7 U.S.C. 2421). The int		421). The information is held
EXHIBIT E STATEMENT OF THE BASIS OF OWNERSHIP	confidential until the certificate is issu	ied (7.0,8.C. 2420).
1. NAME OF APPLICANT(S)	2. TEMPORARY DESIGNATION	3. VARIETY NAME
Texas Agricultural Experiment Station	OR EXPERIMENTAL NUMBER	
	Tx977053	Tamrun OL 02
4. ADDRESS (Street and No., or R.F.D. Ho., City, State, and Z(P) and Country)	5. TELEPHONE (hodow) area code)	5. FAX (Irusulle area node)
Office of the Director, TAES 2147 TAMU	979-845-4747	979-458-4765
College Station, TX 77843-2147	7. PVPO NUMBER	
	1 20030017	7 0
8. Does the applicant own all rights to the variety? Mark an "X" in the section of the configuration of the section of the sec		A
9. Is the applicant (individual or company) a U.S. National or a U.S.	based company? If no, give name of o	country X YES NO
10. Is the applicant the original owner? X YES NO	If no, please answer one of the fo	llowing:
a. If the original rights to variety were owned by individual(s), is	(are) the original owner(s) a U.S. Nation	nal(s)?
YES NO	If no, give name of country	
b. If the original rights to variety were owned by a company(ies	s), is (are) the original owner(s) a U.S. ba	ased company?
YES NO	O If no, give name of country	
11. Additional explanation on ownership (If needed, use the reverse	for extra space):	
TAES policy and handbook manual provide by its employees in the course of their is provided for your records.	<u> </u>	
PLEASE NOTE:		
Plant variety protection can only be afforded to the owners (not licer	nsees) who meet the following criteria:	
 if the rights to the variety are owned by the original breeder, that national of a country which affords similar protection to nationals 	person must be a U.S. national, national of the U.S. for the same genus and spec	of a UPOV member country, or iles.
If the rights to the variety are owned by the company which emplo nationals of a UPOV member country, or owned by nationals of a genus and species.	oyed the original breeder(s), the compan country which affords similar protection	y must be U.S. based, owned by to nationals of the U.S. for the same
3. If the applicant is an owner who is not the original owner, both the	e original owner and the applicant must r	neet one of the above criteria.
The original breeder/owner may be the individual or company who dat for definitions.	firected the final breeding. See Section	41(a)(2) of the Plant Variety Protection
According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponso control number. The valid OMB control number for this information collection is 0581-0055 response, including the time for reviewing the instructions, searching existing data sources	The time required to complete this information colle	ection is estimated to average 6 minutes per

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, sexual orientation, or marital or family status. (Not all prohibited bases apply to all programs). Persons with disabilities who require alternative means for communication of program information (braille, large print, audiotape, etc.) should contact the USDA's TARGET Center at 202-720-2600 (voice and TDD). To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202) 720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.